

(19)



Europäisches Patentamt
European Patent Office
Office européen des brevets

(11) Publication number:

0 343 643
A2

(12)

EUROPEAN PATENT APPLICATION

(21) Application number: 89109406.2

(22) Date of filing: 24.05.89

(51) Int. Cl.4: **A61K 31/41** , **A61K 31/415** ,
A61K 31/42 , **A61K 31/425** ,
A61K 31/44 , **A61K 31/33** ;
A61K 45/06 , **C07D 233/66** ,
C07D 277/32 , **C07D 263/34** ,
C07D 417/06

Claims for the following Contracting States: ES
+ GR

(30) Priority: 25.05.88 US 198528
10.04.89 US 334346

(43) Date of publication of application:
29.11.89 Bulletin 89/48

(84) Designated Contracting States:
AT BE CH DE ES FR GB GR IT LI LU NL SE

(71) Applicant: **WARNER-LAMBERT COMPANY**
201 Tabor Road
Morris Plains New Jersey 07950(US)

(72) Inventor: **Cetenko, Wlaczslaw Antin**
3565 Green Brier Road

Ann Arbor Michigan 48105(US)
Inventor: **Connor, David Thomas**
2453 Antietam

Ann Arbor Michigan 48105(US)
Inventor: **Sorenson, Roderick Joseph**
2820 Briarcliff


Ann Arbor Michigan 48105(US)
Inventor: **Unangst, Paul Charles**
3659 Middleton Drive

Ann Arbor Michigan 48105(US)
Inventor: **Stabler, Stephen Russell**
2211 Latham Street No. 304
Mountainview California 94040(US)

(74) Representative: **Mansmann, Ivo**
c/o Gödecke AG - Patentabteilung Postfach
569 - Mooswaldallee 1-9 —
D-7800 Freiburg(DE)

(54) **Known and selected novel arylmethylenyl derivatives of thiazolidinones, imidazolidinones and oxazolidinones useful as anti-allergy agents and anti-inflammatory agents.**

(57) The present invention is for selected novel compounds, as well as, pharmaceutical compositions and methods of use for known and the selected novel compounds both of the formula



having activity useful for treating allergies and inflammation.


BEST AVAILABLE COPY

EP 0 343 643 A2

KNOWN AND SELECTED NOVEL ARYLMETHYLENYL DERIVATIVES OF THIAZOLIDINONES, IMIDAZOLIDINONES AND OXAZOLIDINONES USEFUL AS ANTIALLERGY AGENTS AND ANTIINFLAMMATORY AGENTS

It is now found that compounds from among known and selected novel arylmethylenyl derivatives of thiazolidinones, oxazolidinones or imidazolidinones have activity useful for treating allergies or inflammation.

Thus, the present invention is for selected novel compounds, as well as, pharmaceutical compositions and methods of use for known compounds and the selected novel compounds of the formula (I)



I

and pharmaceutically acceptable salts thereof; wherein Ar is (i) phenyl unsubstituted, (ii) phenyl substituted by from one to three of lower alkyl, lower alkoxy, hydroxy, halogen, trifluoromethyl, $\text{NR}_{10}\text{R}_{11}$ wherein R_{10} and R_{11} are independently hydrogen or lower alkyl, NO_2 , mercapto, or lower alkylthio, (iii) naphthyl; (iv) benzofuranyl, (v) benzothiophenyl, (vi) 2- or 3-thienyl, (vii) 2- or 3-indolyl, (viii) 2- or 3-furanyl, or (ix) 2-, 3-, or 4-pyridyl

Y and Y_1 is oxygen or sulfur;


X is sulfur, oxygen, NH or NCH_3 and

X_1 is NH or NCH_3

with the proviso that when Ar is substituted phenyl, the substituents are either not 3,5-di(isopropyl)-4-hydroxy or if $\text{X}_1 = \text{NH}$, $\text{X} = \text{S}$, $\text{Y} = \text{S}$ not 3,5-di(t-butyl) together with 4-hydroxy.

Japanese Application No. 84-133826 as found in Derwent Abstracts No. 87-076379/11 discloses new 3,5-diisopropylbenzylidene-2,4-imidazolidinone and the analogous thiazoles with antiallergic and tyrosine kinase inhibiting activity as well as other heterocyclic containing compounds. EP No. 211,670A as described in Derwent 87-051809/08 describes compounds excluded by the proviso above having a 3,5-diisobutylbenzylidene substituent useful for treating inflammation, ischaemia induced cell damage and arthritis. Thus, known heterocyclic compounds are not as closely related as these named here and are excluded from the present invention.

A French Patent No. 2,169,334, C. H. Boehringer Sohn, October 12, 1973 describes a series of N-substituted oxazoles as antiarthritics, antirheumatics, and immunosuppressants having the formula



which are not included in the present invention.

Copending application United States Ser.No. 277,171 is for novel compounds having a 3,5-di-tertiarybutyl-4-hydroxyphenyl moiety so its disclosure relating to references in the Background of the Invention is incorporated herein by reference.

The compounds of the formula I have a hitherto unknown pharmaceutical activity useful for treating allergic or inflammatory conditions of diseases. The invention compounds are now also found to have activity as inhibitors of 5-lipoxygenase and/or cyclooxygenase providing treatment of conditions advantageously affected by such inhibition including inflammation, arthritis, pain, pyrrhia, and the like. Thus, the present invention is also a pharmaceutical composition for diseases or conditions advantageously affected by activity inhibiting singly or together 5-lipoxygenase and cyclooxygenase or method of use therefor.

Therefore, the present invention is particularly a pharmaceutical composition for the treatment of allergy or inflammation which comprises an antiallergy or antiinflammatory effective amount of a compound of the formula I together with a pharmaceutically acceptable carrier. Further, the present invention is the use of the above compounds for the preparation of pharmaceuticals for treating allergies or inflammation in subject

suffering therefrom.

The present invention is also the novel selected compounds of the formula I noted hereinafter having activity heretofore unknown for analogous compounds.

Among the known compounds for the compositions or methods of use of the present invention the following are references for compounds listed for the Examples of the invention:

<u>Example No.</u>	<u>Reference</u>
1	J. Chem. Soc., pp. 958-65 (1946)
2	J. Org. Chem., <u>26</u> , p. 1326 (1961)
3	Chem. Ber., pp. 459-69 (1941)
4	J. Chem. Soc., pp. 3547-52 (1950)
5	J.C.S., p. 1165 (1954)
6	J. Org. Chem., p. 1326 (1961)
7	U.S. Pat. No. 3,843,797/JACS, p. 2357 (1951)

<u>Example No.</u>	<u>Reference</u>
8	C.A. <u>89</u> : 100767m, JACS, p. 2357 (1951)
9	Tet., p. 2781 (1969)
10	Canadian J. Chem., p. 2089 (1959)
12	Ger. Pat. DE 3,433,475/C.A. <u>103</u> , 928404
17	J.O.C., p. 32 (1956)
18	Tet., p. 2781 (1969)
19	Tet., p. 2781 (1969)
20	Tet., p. 2781 (1969)
21	J. Ind. Chem. Soc., p. 77 (1984)
22	J. Ind. Chem. Soc., p. 77 (1984)
26	See Example 8 above
28	Int. J. Sulf. Chem. Part A, p. 261 (1972)
29	C.A. <u>76</u> (5), 25152h
30	C.A. <u>79</u> (25), 143522p
32	C.A. <u>76</u> (5), 25152h
34	C.A. <u>79</u> (25), 143522p
36	Biol. Mem. <u>9</u> , 200 (1984)
38	Zh. Org. Khim, p. 212 (1983)
40	J.C.S. (1), p. 385 (1984)
41	C.A. <u>89</u> : 100767n
42	Chem. Phar. Bull., p. 1619 (1986)
43	Int. J. Sulf. Chem. Part A, p. 261 (1972)
44	J. Pharm. Soc. Jap., p. 154 (1956)
45	Ger. Pat. DE 3,433,475
48	J.C.S., p. 3547 (1950)
49	C.A. <u>47</u> , 9543a
50	C.A. <u>47</u> , 9543a
68	C.A. <u>71</u> (17), 81253a
69	Zh. Ob. Khim <u>26</u> , p. 3092 (1956)
71	Ukrain Khim Zh. <u>16</u> , p. 545 (1950)
83	Chem. Pharm. Bull., p. 1619 (1986)
86	Am. Chem. J., p. 368 (1911)
87	JACS, p. 1606 (1913)
90	JACS, p. 1606 (1913)
91	Monats, p. 352 (1961)
92	Chem. Pharm. Bull., p. 1619 (1986)

<u>Example No.</u>	<u>Reference</u>
96	Roecz, Chem., p. 1381 (1984)
103	U.S. Pat. No. 3,017,270
104	Egypt J. Chem., <u>26</u> , 301 (1983)
105	French Pat. No. 1,604,530

The selected novel compounds of the present invention are found in Examples 11, 13-16, 51-53, 53-63, 64, 70, 72-79, 80, 82, 88, 97, and 99-101, 106-111.


Accordingly, the present invention is

- (a) a selected novel compound as in the noted examples heretofor or a salt thereof;
- (b) a method for preparing a novel selected compound or a pharmacologically acceptable salt thereof;
- (c) a pharmaceutical composition comprising a compound of formula (I) or a physiologically acceptable salt thereof and a pharmaceutically acceptable carrier therefor;
- (d) a method for preparing such compositions;
- (e) a method for the inhibition of histamine by use of a nontoxic, effective, inhibitory amount of a compound of formula I or a physiologically acceptable salt thereof;
- (f) a method for the prophylaxis or treatment of disease or condition in a mammal, including man, comprising the administration to said mammal of a nontoxic, therapeutically or prophylactically effective amount of a compound of formula I or a physiologically acceptable salt thereof;
- (g) a method for the prophylaxis or treatment of any individual condition described herein, in a mammal, including man, comprising the administration to said mammal of a nontoxic therapeutically or prophylactically effective amount of a compound of formula I or a physiologically acceptable salt thereof;
- (h) a method for the prophylaxis or treatment of allergy or inflammatory condition in a mammal, including man, comprising administration to said mammal of a nontoxic, effective, antiallergic or antiinflammatory amount of a compound of formula I or a physiologically acceptable salt thereof;
- (i) a novel compound as set out in the examples noted heretofor or a physiologically acceptable salt thereof for use in medicine, especially as defined in (f)-(h) above;
- (j) use of a compound of formula I or a physiologically acceptable salt thereof in the manufacture of medical therapeutic agents, particularly those for use as defined in (f)-(h) above; and
- (k) any novel feature described herein.

SUMMARY OF THE INVENTION

The present invention is a pharmaceutical composition and method of use as defined above for compounds of the formula I as well as the novel compounds also as noted above.


A preferred embodiment of the present invention is the composition or method of use for a compound of the formula (II)



or pharmaceutically acceptable salts thereof; wherein X and Y are as defined above, R_1 , R_2 , R_3 , R_4 and R_5 are independently H, lower alkyl, lower alkoxy, OH, halogen, CF_3 , $NR_{10}R_{11}$ wherein R_{10} and R_{11} are as defined above, NO_2 , mercapto, or lower alkylthio, wherein of from one to three of R_1 , R_2 , R_3 , R_4 , and R_5

are other than hydrogen, but excluding R_1 , R_2 , R_3 , R_4 and R_5 which are 3,5-di(2-isopropyl) and 4-OH or which are 3,5-di(t-butyl) and 4-OH when X is NH, or both X and Y are S.


A preferred embodiment of the formula II of the present invention is the composition or method of use for a compound of the formula (III)



III

or pharmaceutically acceptable salts thereof; wherein R_1 , R_2 , R_3 , R_4 and R_5 are as defined above.


Another preferred embodiment of the formula II of the present invention is the composition or method of use for a compound of the formula (IV)



IV

or pharmaceutically acceptable salts thereof; wherein R_1 , R_2 , R_3 , R_4 and R_5 are as defined above.


Also among the preferred embodiment of the formula II is the composition and method of use for the compound of the formula (V)



V

or pharmaceutically acceptable salts thereof; wherein R_1 , R_2 , R_3 , R_4 and R_5 are as defined above.


Another preferred embodiment of the formula II is the composition and method of use for the compound of the formula (VI)



VI

or pharmaceutically acceptable salts thereof; wherein R_1 , R_2 , R_3 , R_4 and R_5 are as defined above.


Another preferred embodiment of the formula II is the composition and method of use for the compounds of the formula (VII)



VII


or pharmaceutically acceptable salts thereof; wherein X is NH or NCH₃ and R₁, R₂, R₃, R₄ and R₅ is as defined above.

Another preferred embodiment of the formula II is the composition and method of use for the compounds of the formula (VIII)



VIII

or pharmaceutically acceptable salts thereof wherein R₁, R₂, R₃, R₄, and R₅ are as defined above. And finally, a preferred embodiment of the formula II is the composition and methods of use for the compounds of the formula VIIIA



VIIIA

or pharmaceutically acceptable salts thereof; wherein R₁, R₂, R₃, R₄ and R₅ are as defined above.

The most preferred embodiment of the formula II is the composition and method of use for the following compounds which inhibit the release of histamine from human basophils as described hereinafter in the HHB assay stimulated with anti IgE at an IC₅₀ less than 20 μM.

- 5-[(2,5-dimethoxyphenyl)methylene]-2-thioxo-4-thiazolidinone,
- 5-[2,4,5-trimethoxyphenyl)methylene]-2-thioxo-4-thiazolidinone,
- 5-[(3,4,5-trimethoxyphenyl)methylene]-2-thioxo-4-thiazolidinone,
- 5-[(3,4-dichlorophenyl)methylene]-2-thioxo-4-thiazolidinone,
- 5-[(3,4-dimethoxyphenyl)methylene]-2-thioxo-4-thiazolidinone,
- 5-[(3,5-dimethoxy-4-hydroxyphenyl)methylene]-2-thioxo-4-thiazolidinone,
- 5-[(3,5-dimethyl-4-hydroxyphenyl)methylene]-2-thioxo-4-thiazolidinone,
- 5-[(5-bromo-4-hydroxy-3-methoxyphenyl)methylene]-2-thioxo-4-thiazolidinone,
- 5-[(4-methoxyphenyl)methylene]-2-thioxo-4-thiazolidinone,
- 5-[(5-hydroxy-4-methoxyphenyl)methylene]-2-thioxo-4-thiazolidinone,
- 5-[(3,5-dimethoxyphenyl)methylene]-2-thioxo-4-thiazolidinone,
- 5-[(4-hydroxy-3,5-dimethoxyphenyl)methylene]-2,4-thiazolidinedione,
- 5-[(4-hydroxy-3-methoxyphenyl)methylene]-2,4-thiazolidinedione,
- 5-[(4-hydroxy-3-methoxyphenyl)methylene]-2-thioxo-4-oxazolidinone,
- 5-[(4-hydroxy-3,5-dimethoxyphenyl)methylene]-2-thioxo-4-oxazolidinone,
- 2-thioxo-5-[(3,4,5-trimethoxyphenyl)methylene]-4-oxazolidinone,
- 5-[(4-hydroxy-3,5-dimethoxyphenyl)methylene]-2-thioxo-4-imidazolidinone.

Novel compounds which inhibit singly or together 5-lipoxygenase and cyclooxygenase are found by determining percent inhibition in ARBL/ARBC Whole Cell 5-lipoxygenase and Cyclooxygenase Assays and Carrageenan-Induced Rat Foot Paw Edema-2 (CFE-2) Assay described hereinafter.

Among the preferred compounds as determined in the ARBL/ARBC and CFE-2 assays are the

following:

- 2,4-thiazolidione, 5-[[3,5-bis(1,1-dimethylethyl)-4-hydroxyphenyl]methylene]-, (E)-
 4-oxazolidinone, 5-[[3,5-bis(1,1-dimethylethyl)-4-hydroxyphenyl]methylene]-2-thio-
 2,4-oxazolidinone, 5-[[3,5-bis(1,1-dimethylethyl)-4-hydroxyphenyl]methylene]-,
 5 2,4-oxazolidinedione, 5-[(4-hydroxy-3,5-dimethoxyphenyl)methylene]-, and
 4-thiazolidinone, 5-[(4-bromophenyl)methylene]-2-thio-.

The most preferred novel compounds of the present invention now found having activity which inhibits 5-lipoxygenase and cyclooxygenase are 5-[[3,5-bis(1,1-dimethylethyl)-4-hydroxyphenyl]methylene]-2,4-thiazolidinedione and 5-[(3,5-bis(1,1-dimethylethyl)-4-hydroxyphenyl)-2-thio-4-oxazolidinone.

- 10 Thus, the present invention is also a pharmaceutical composition for the treatment of conditions advantageously affected by the inhibition of 5-lipoxygenase and/or cyclooxygenase which comprises an amount effective for the treatment of the condition of a compound of the formula I or various preferred embodiments and the pharmaceutically acceptable acid addition or base salt thereof together with a pharmaceutically acceptable carrier. The condition is meant to include, for example, arthritis or other
 15 inflammatory diseases, allergic conditions or diseases, pain, fever, and psoriasis, but now preferably inflammatory diseases.

- The present invention is also a method for treatment of the condition as noted above in a mammal, including humans, suffering therefrom with a compound of the formula I or the pharmaceutically acceptable acid addition or base salt thereof, in unit dosage form. The invention also provides for use of any such
 20 compound of formula I or particularly various preferred embodiments or salt thereof in the manufacture of medical therapeutic agent.

Pharmaceutical composition or use of the compound or salt of formula I is meant to include treatment understood to be prophylactic pertinent to the foregoing named condition.

25

A DETAILED DESCRIPTION OF THE INVENTION

- In the present invention "lower alkyl" is alkyl of from one to six carbons, inclusive, and means methyl,
 30 ethyl, propyl, butyl, pentyl, or hexyl and isomers thereof.

"Lower alkoxy" means methoxy, ethoxy, propoxy, butoxy, pentoxy, or hexoxy and isomers thereof.

"Lower alkylthio" means methylmercapto, ethylmercapto, propylmercapto, butylmercapto, pentylmercapto, or hexylmercapto and isomers thereof.

"Halogen" is chloro, bromo, fluoro, or iodo.

- 35 Generally, the selected novel compounds of formula I as well as the known compounds are prepared by processes that are known or are prepared by processes analogous to those that are known from known starting materials or starting materials that can be prepared by known methods. For example, the following starting materials are obtained as follows:

- 3-Bromo-4-hydroxybenzaldehyde and 3,5-dibromo-4-hydroxybenzaldehyde are prepared according to Paal,
 40 Chem. Ber. 28 (1895), 2407.

4,5-Dimethoxy-2-hydroxybenzaldehyde is prepared according to Robinson & Head, J. Chem. Soc. (1930), 2440.

3-Methoxy-5-hydroxybenzaldehyde is prepared according to Ben, et al, J. Org. Chem. 50 (1985), 2238.

- 3,5-Diisopropyl-4-hydroxybenzaldehyde and 3,5-dimethyl 4-hydroxybenzaldehyde are prepared according
 45 to U.S. Patent No. 4,009,210.


3-(dimethylamino)methyl-4-hydroxy-5-methoxybenzaldehyde is prepared according to Hemetsberger, Mon-ats. Chem. 102 (1971), 1110.

3-Methylhydantoin is prepared according to Güler and Moodie, J. Chem. Soc. Perk II (1980), 1752.

A scheme for preparation of the compounds of formula I above is as follows:

50


55



Scheme 1

wherein Ar, X₁, X and Y are as defined above.


More particularly, the present invention includes compounds of type 1:



where R₁, R₂, R₃, R₄ and R₅ are as defined above. These compounds are named as substituted 5-phenylmethylene-2-thioxo-4-oxazolidinones, or 5-phenylmethylene-2,4-oxazolidinediones.

Salts of these compounds can be formed by treatment with organic and inorganic bases due to the presence of an acidic NH.

Certain thiazolidinones and oxazolidinones are acidic (pK_a ~ 3-6) due to the presence of a tautomeric proton on the heterocyclic ring:




These compounds can form salts with inorganic and organic bases, including choline derivatives.

If the parent oxazolidinone or thiazolidinone and choline free base or a choline derivative (such as choline bicarbonate or choline chloride) are combined in a suitable solvent (such as an alcohol, water, or an alcohol/water solution), the choline salt can generally be obtained by precipitation from solution or by evaporation of the reaction solution.

A general method of preparation of these compounds is the aldol condensation of an aldehyde 2 with an active methylene compound 3 (Scheme 1 above). This condensation can be carried out in alcoholic solvents in the presence of a base such as ammonia, ammonium salts, or piperidine, or with mineral acid

Scheme 1'

2

I wherein X is O

X₁ is NH

Y is S

15

20

catalysis. A particularly favored procedure is the use of anhydrous sodium acetate in glacial acetic acid, with heating at reflux for 1-24 hours. References to this procedure include: G. R. Newkome and A. Nayak, in *Advances in Heterocyclic Chemistry*, A. R. Katritzky and A. J. Boulton, Eds., Academic Press, New York, NY, Vol. 25, p. 83 and N. K. Ushenko and T. E. Gorizdra, *Ukrain Khim. Zhur.*, 16, 545 (1950).

25

An alternate procedure for the preparation of the above compound when Y = O is the preparation of the corresponding oxazole 4 followed by oxidation/hydrolysis to the desired oxazole I wherein X is O, X₂ is NH and Y is O (Scheme 2):

Scheme 2

4

I wherein X is O,

X₁ is NH, and

Y is O.

40


45

For the preparation of 4, see, for example, F. C. Brown, *Chem. Rev.*, 61, 463 (1961), and the (Newkome) reference previously cited. For the conversion 4 to I of Scheme 2, see also J. W. Clark-Lewis, *Chem. Rev.*, 58, 63 (1958), and N. A. Shenberg, L. S. Guseva, A. I. Ginak, and E. G. Sochilin, *Zhur. Organ. Khim.*, 14, 1323 (1978).

50


Variations of the above Scheme 1 are also readily prepared by the methods of Scheme 1' discussed above. These are shown here in Scheme 1" as follows.

Scheme 1"



20 An alternate procedure for the preparation of the above noted oxazole I wherein X is O, X₁ is NH, and Y is O is by an alkylation/hydrolysis procedure (Scheme 2') as follows:

Scheme 2'



45 The procedure of Scheme 2' is essentially that of N. A. Shenberg, L. S. Guseva, A. I. Ginak, and E. G. Sochilin, Zhur. Organ. Khim., 14, 1323 (1978); and J. S. H. Davies, W. Hook, and F. Long, J. Chem. Soc., 30 (1950).

50 The compounds of formula I are useful both in the free acid form, in the form of base salts where possible, and in the form of acid addition salts. The three forms are within the scope of the invention. In practice, use of the salt form amounts to use of the base form. Appropriate pharmaceutically acceptable salts within the scope of the invention are those derived from mineral acids such as hydrochloric acid and sulfuric acid; and organic acids such as methanesulfonic acid, benzenesulfonic acid, p-toluenesulfonic acid, and the like, giving the hydrochloride, sulfamate, methanesulfonate, benzenesulfonate, p-toluenesulfonate, and the like, respectively or those derived from bases such as suitable organic and inorganic bases. Examples of suitable inorganic bases for the formation of salts of compounds of this invention include the hydroxides, carbonates, and bicarbonates of ammonia, sodium, lithium, potassium, calcium, magnesium, 55 aluminum, zinc, and the like.

Salts may also be formed with suitable organic bases. Bases suitable for the formation of pharmaceutically acceptable base addition salts with compounds of the present invention include organic bases which are nontoxic and strong enough to form such salts. These organic bases form a class whose limits are

readily understood by those skilled in the art. Merely for purposes of illustration, the class may be said to include mono-, di-, and trialkylamines, such as methylamine, dimethylamine, and triethylamine; mono-, di-, or trihydroxyalkylamines such as mono-, di-, and triethanolamine; amino acids such as arginine, and lysine; guanidine; N-methylglucosamine; N-methylpiperazine; morpholine; ethylenediamine; N-benzyl-phenethylamine; tris(hydroxymethyl) aminomethane; and the like. (See for example, "Pharmaceutical Salts," J. Pharm. Sci., 66(1), 1-19 (1977).)

The acid addition salts of said basic compounds are prepared either by dissolving the free base of compound I in aqueous or aqueous alcohol solution or other suitable solvents containing the appropriate acid or base and isolating the salt by evaporating the solution, or by reacting the free base of compound I with an acid as well as reacting compound I having an acid group thereon with a base such that the reactions are in an organic solvent, in which case the salt separates directly or can be obtained by concentration of the solution.

The base salts of compounds of formula I described above are prepared by reacting the appropriate base with a stoichiometric equivalent of the acid compounds of formula I to obtain pharmacologically acceptable base salts thereof.

The acid solution salts of said basic compounds are prepared either by dissolving the free base in aqueous or aqueous alcohol solution or other suitable solvents containing the appropriate acid and isolating the salt by evaporating the solution, or by reacting the free base and acid in an organic solvent, in which case the salt separates directly or can be obtained by concentration of the solution.

The compounds of this invention may also exist in hydrated or solvated forms.

The products of the reactions described herein are isolated by conventional means such as extraction, distillation, chromatography and the like.

The antiallergic and antiinflammatory activity of the compounds having the formula I of the present invention is determined by an assay showing inhibition of the release of histamine from human basophils (HNB). A description of the protocol of the HNB assay is found hereinafter.

Thus, pharmaceutical compositions are prepared from the compounds of formula I and salts thereof described as the present invention in unit dosage form comprising the compound either alone or in admixture with a pharmaceutically acceptable carrier appropriately selected from those known.

A physician or veterinarian of ordinary skill readily determines a subject who is exhibiting allergic or inflammatory symptoms. Regardless of the route of administration selected, the compounds of the present invention are formulated into pharmaceutically acceptable dosage forms by conventional methods known to the pharmaceutical art.

The compounds can be administered in such oral unit dosage forms such as tablets, capsules, pills, powders, or granules. They also may be administered rectally or vaginally in such forms as suppositories or bougies; they may also be introduced parenterally (e.g., subcutaneously, intravenously, or intramuscularly), using forms known to the pharmaceutical art. They are also introduced directly to an affected area (e.g., in the form of eye drops or by inhalation). For the treatment of allergic or inflammatory conditions such as erythema, the compounds of the present invention may also be administered topically in the form of ointments, creams, gels, or the like. In general, the preferred route of administration is orally.

An effective but nontoxic quantity of the compound I is employed in treatment. The ordinarily skilled physician or veterinarian will readily determine and prescribe the effective amount of the antiallergic or antiinflammatory agent to prevent or arrest the progress of the condition. The dosage regimen is selected in accordance with a variety of factors including the type, age, weight, sex, and medical condition of the mammal, the severity of symptoms of the disease being treated, the route of administration and particular compound of formula I employed. An ordinarily skilled physician or veterinarian will readily determine and prescribe the effective amount of the compound I to prevent or arrest the progress of the condition. In so proceeding, the physician or veterinarian could employ relatively low dosages at first, subsequently increasing the dose until a maximum response is obtained. For convenience, the total daily dosage may be divided and administered in portions during the day if desired. In so proceeding, the physician or veterinarian could employ relatively low dosages at first, subsequently increasing the dose until a maximum response is obtained.

Initial dosage of the compounds of the invention having formula I are ordinarily in the area of 10 mg up to 2 g per day orally, preferably 20 mg to 500 mg per dose orally, given from one to four times daily or as needed. When other forms of administration are employed, equivalent doses are administered.

A suitable dose of a compound of formula (I) or pharmacologically acceptable salt thereof for a mammal suffering from, or likely to suffer from any condition as described hereinbefore is 0.1 μ g - 500 mg of the compound per kilogram body weight. In the case of systemic administration, the dose may be in the range of 0.5 to 500 mg of the compound per kilogram body weight, the most preferred dosage being 0.5 to 50

mg/kg of mammal body weight administered two or three times daily. In the case of topical administration, e.g., to the skin or eye, a suitable dose may be in the range 0.1 ng-100 µg of the compound per kilogram, typically about 0.1 µg/kg.

In the case of oral dosing for the treatment or prophylaxis of arthritis or inflammation in general, due to any course, a suitable dose of a compound of formula (I) or physiologically acceptable salt thereof, may be as specified in the preceding paragraph, but most preferably is from 1 mg to 10 mg of the compound per kilogram, the most preferred dosage being from 1 mg to 5 mg/kg of mammal body weight, for example, from 1 to 2 mg/kg.

It is understood that the compositions and methods of treatment of the present invention as described above also include the free acid, the pharmacologically acceptable base salts and acid addition salts of the compounds of formula I.

The following Examples further illustrate the invention, but are not meant to be limiting thereto.


EXAMPLE 1

5-[(2,3-Dimethoxyphenyl)methylene]-2-thioxo-4-thiazolidinone

A mixture of 2,3-dimethoxybenzaldehyde (5.2 g, 30 mmoles), rhodanine (4.0 g, 29 mmoles), sodium acetate (8.4 g, 102 mmoles), and acetic acid (50 ml) is stirred under an inert atmosphere and heated to reflux. After 4 hours the mixture is stirred into water (250 ml) and the precipitate is filtered off, rinsed successively with water (3X), ethanol (2X), and ether (2X), and dried to afford the pure product (7.8 g), mp 268-269° C.

TABLE 1

The following examples are prepared from the corresponding benzaldehydes using the procedure described in Example 1.



5

10

Example	R ₁	R ₂	R ₃	R ₄	mp °C
2	OMe	H	OMe	H	275-276
3	OMe	H	H	OMe	245-246
4	OMe	OMe	OMe	H	205-206
5	OMe	H	OMe	OMe	297 (dec)
6	H	OMe	OMe	OMe	200-201
7	Cl	H	Cl	H	234-235 (Recryst. from DMF)
8	H	Cl	Cl	H	241-242
9	H	OMe	OMe	H	232-234
10	H	OMe	OH	OMe	252-253 (Recryst. from MeOH)
11	H	F	OMe	H	264-265
12	H	OH	OH	H	>300
13	H	Br	OH	H	283-285 (Recryst. from EtOH)
14	H	Br	OH	Br	>300
15	H	Me	OH	Me	293-295
16	H	OMe	OH	Br	275 (dec)
17	H	H	H	H	204-205
18	MeO	H	H	H	249-250
19	H	MeO	H	H	232-233
20	H	H	MeO	H	250-251
21	H	OH	H	H	246-248
22	H	H	OH	H	288 (dec)
23	H	OEt	H	H	170-172
24	H	OCHMe ₂	H	H	191-193
25	H	OCH ₂ Ph	H	H	197-198
26	H	Me	H	H	198-200
27	H	CF ₃	H	H	189-190
28	H	Cl	H	H	233-235

55

TABLE I (cont'd)

	Example	R ₁	R ₂	R ₃	R ₄	mp °C
5	29	H	Br	H	H	239-241
	30	H	NO ₂	H	H	265-267
	31	H	H	Cl	H	224-226
10	32	H	H	Br	H	236-238
	33	H	H	OPh	H	150-151
	34	H	H	NO ₂	H	265-267
	35	H	H	SMe	H	258-260
15	36	H	MeO	OH	H	229-230
	37	H	MeO	H	MeO	257-259
	38	H	H	Me	H	225-227
20	39	H	H	Ph	H	240-242

EXAMPLE 40

5-[(2,3-Dichlorophenyl)methylene]-2-thioxo-4-thiazolidinone

A mixture of 2,3-dichlorobenzaldehyde (4.6 g, 26 mmoles), rhodanine (3.5 g, 25 mmoles), sodium acetate (7.5 g, 91 mmoles), and acetic acid (40 ml) is stirred under an inert atmosphere and heated to reflux. After 2.5 hours the mixture is stirred into ice-water (250 ml) and the precipitate is filtered off, rinsed three times with water and dried. Recrystallization from methanol gave the pure product (2.9 g), mp 203-204°C.

The following examples are prepared from the corresponding benzaldehydes using the procedure described in Example 40:

	Example	R ₁	R ₂	R ₃	R ₄	R ₅	mp °C	Recryst. Solvent
40	41	Cl	H	H	H	Cl	179-181	---
	42	H	tBu	OH	tBu	H	248-249	EtOAc
	43	H	F	H	H	H	201-202	MeCN
45	44	OH	H	H	H	H	216 (dec)	MeCN
	45	H	OH	OMe	H	H	224-225	MeOH
	46	H	iPr	OH	iPr	H	200-202	MeCN
	47	H	OH	OMe	OMe	H	212-213	MeOH
	48	OMe	H	H	H	OMe	273-275	DMF
50	49	OH	H	H	OMe	H	253 (dec)	---
	50	H	OH	H	OMe	H	259-260	THF
	51	H	OMe	OH	tBu	H	233-234	MeCN
	52	OH	H	OMe	OMe	H	258 (dec)	---

EXAMPLE 53

5-[[3-[(Dimethylamino)methyl]-4-hydroxy-5-methoxyphenyl]methylene]-2-thioxo-4-thiazolidinone

A mixture of 3-[(dimethylamino)methyl]-4-hydroxy-5-methoxybenzaldehyde (2.5 g, 12 mmoles), rhodanine (1.6 g, 12 mmoles), β -alanine (0.6 g, 7 mmoles), and acetic acid (35 ml) is stirred under an inert atmosphere and heated to reflux. After 4 hours the acetic acid is removed by rotary evaporator and the residue dissolved in water (250 ml). The solution is brought to approximate neutrality by addition of saturated aqueous sodium bicarbonate, and stirred for 3 hours. The precipitate is filtered off, rinsed three times with water and dried, then triturated in boiling methanol (100 ml), cooled, filtered off and dried to afford the pure product (2.9 g), mp 198° C (dec).

EXAMPLE 54

5-[(5-Bromo-2-thienyl)methylene]-2-thioxo-4-thiazolidinone

Prepared by the method described in Example 1 from 5-bromo-2-thiophenecarboxaldehyde, mp 250-252° (dec).

INTERMEDIATE A

N,6-Dimethoxy-N-methyl-2-benzofurancarboxamide

To a stirred solution of 6.0 g (0.031 mole) of 6-methoxy-2-benzofurancarboxylic acid (A. McGookin, A. Robertson, and W. B. Whalley, J. Chem. Soc. 787 (1940)) in 100 ml of tetrahydrofuran (under a nitrogen atmosphere) was added in one portion, 3.3 ml (4.8 g; 0.038 mole) of oxalyl chloride. The mixture was stirred for 5 minutes, then treated with 3 drops of N,N-dimethylformamide. After stirring at room temperature for 2 hours, the solvent was evaporated to leave a residue of crude 6-methoxy-2-benzofurancarbonyl chloride, mp 96-99° (a mp of 101° is given by A. McGookin, etc., cited above). The residue was re-dissolved in 100 ml of dichloromethane, and the solution was filtered. The filtrate was added dropwise over 20 minutes to an ice-cooled solution of 3.6 g (0.037 mole) of N,O-dimethylhydroxylamine hydrochloride and 9.4 ml (7.7 g; 0.077 mole) of 1-methylpiperidine in 100 ml of dichloromethane. After stirring for 2 hours, the mixture was added to 400 ml of ice-cold 5% aqueous hydrochloric acid. The layers were separated, and the aqueous layer was extracted with fresh dichloromethane (3X 100 ml). The combined organic layers were washed with water (1X 200 ml), 3% aqueous sodium bicarbonate (2X 200 ml), and water again. The extracts were dried (anhydrous magnesium sulfate) and evaporated. Trituration of the residue with ether yielded 5.4 g (74% yield) of the analytically pure amide product, mp 73-77°.

Calcd. for C ₁₂ H ₁₃ NO ₄ :	C, 61.27;	H, 5.57;	N, 5.96
Found:	C, 61.26;	H, 5.52;	N, 5.78.

INTERMEDIATE B

6-Methoxy-2-benzofurancarboxaldehyde

An ice-cooled solution of 4.0 g (0.017 mole) of N,6-dimethoxy-N-methyl-2-benzofurancarboxamide in

100 ml of tetrahydrofuran was treated cautiously over 15 minutes (under a nitrogen atmosphere) with 0.65 g (0.017 mole) of lithium aluminum hydride. The mixture was stirred for 1 hour, then quenched by the careful addition of 100 ml of saturated aqueous sodium bisulfate solution. The reaction mixture was extracted with ether (3X 75 ml), and the combined extracts were washed with cold 2% aqueous hydrochloric acid (2X 100 ml), followed by brine (1X 100 ml). The organic layer was dried (anhydrous magnesium sulfate) and evaporated. The residue was chromatographed (silica gel, dichloromethane elution) to yield 2.2 g (74% yield) of the analytically pure aldehyde, mp 75-77°.

Calcd. for $C_{10}H_8O_3$:	C, 68.18;	H, 4.58
Found:	C, 68.03;	H, 4.48.

EXAMPLE 55

5-[(6-Methoxy-2-benzofuranyl)methylene]-2-thioxo-4-thiazolidinone

A mixture of 1.5 g (0.0085 mole) of 6-methoxy-2-benzofurancarboxaldehyde, 1.1 g (0.0083 mole) of 2-thioxo-4-thiazolidinone ("rhodanine"), and 2.5 g (0.030 mole) of anhydrous sodium acetate in 40 ml of glacial acetic acid (under a nitrogen atmosphere) was stirred and heated at reflux for 5 hours. The cooled mixture was added to 250 g of ice/water. The precipitated solid was filtered, washed with water, then with a little cold ethanol to yield 2.0 g (81% yield) of the analytically pure thiazole, mp > 290°.

Calcd. for $C_{13}H_9NO_3S_2$:	C, 53.59;	H, 3.11;	N, 4.81;	S, 22.01
Found:	C, 53.82;	H, 3.08;	N, 5.15;	S, 21.71

The following Examples are for compounds also prepared by the procedures described the preparation of intermediates A and B and Example 44 above.

EXAMPLE 56

5-[(5-Methoxy-2-benzofuranyl)methylene]-2-thioxo-4-thiazolidinone

mp 261-263° (recrystallized from acetonitrile/N,N-dimethylformamide).

Calcd. for $C_{13}H_9NO_3S_2$:	C, 53.59;	H, 3.11;	N, 4.81;	S, 22.01
Found:	C, 53.68;	H, 3.23;	N, 4.87;	S, 21.88.

The starting carboxylic acid was 5-methoxy-2-benzofurancarboxylic acid (W. B. Whalley, J. Chem. Soc., 3479 (1953)).

EXAMPLE 57

5-[(Benzo[b]thien-2-yl)methylene]-2-thioxo-4-thiazolidinone

mp 245°-dec (recrystallized from acetonitrile/N,N-dimethylformamide).

Calcd. for $C_{12}H_7NOS_3$:	C, 51.96;	H, 2.54;	N, 5.05;	S, 34.68
Found:	C, 51.87;	H, 2.41;	N, 5.13;	S, 34.45.

5 The starting carboxylic acid was commercially available benzo[b]thiophene-2-carboxylic acid.

EXAMPLE 58

10 5-[[5-Methoxy-3-(1-methylethoxy)benzo[b]thien-2-yl]methylene]-2-thioxo-4-thiazolidinone

mp 242° -dec.

Calcd. for $C_{15}H_{15}NO_3S_3$:	C, 52.58;	H, 4.14;	N, 3.83;	S, 26.32
Found:	C, 53.00;	H, 4.13;	N, 3.75;	S, 25.95

20 The starting carboxylic acid was 3-(1-methylethoxy)-5-methoxy-benzo[b]thiophene-2-carboxylic acid (D. T. Connor, W. A. Cetenko, P. C. Unangst, and E. A. Johnson, U.S. Patent No. 4,703,053.

EXAMPLE 59

25 5-[(5-Methoxy-1-phenyl-1H-indol-2-yl)methylene]-2-thioxo-4-thiazolidinone

30 mp 250-253° (recrystallized from aqueous acetonitrile/N,N-dimethylformamide).

Calcd. for $C_{19}H_{14}N_2O_2S_2 \cdot 1H_2O$:	C, 59.35;	H, 4.19;	N, 7.29;	S, 16.68
Found:	C, 59.56;	H, 4.03;	N, 7.06;	S, 16.02.

35 The starting carboxylic acid was 5-methoxy-1-phenyl-1H-indole-2-carboxylic acid (P. C. Unangst, D. T. Connor, S. R. Stabler, and R. J. Weikert, J. Heterocyclic Chem., 24, 811 (1987)).

EXAMPLE 60

40 5-[(5-Methoxy-1-phenyl-1H-indol-2-yl)methylene]-2-thioxo-4-thiazolidinone, 1-methylpiperidine salt

45 A suspension of 3.0 g (0.0082 mole) of 5-[(5-methoxy-1-phenyl-1H-indol-2-yl)methylene]-2-thioxo-4-thiazolidinone in 50 ml of methanol was treated with 1.1 ml (0.89 g; 0.0090 mole) of 1-methylpiperidine. The mixture was stirred briefly on the steam bath until homogeneous and filtered warm. Cooling to room temperature overnight yielded 3.2 g (84% yield) of the crystalline, analytically pure salt, mp 188-191°.

Calcd. for $C_{19}H_{14}N_2O_2S_2 \cdot C_6H_{13}N$:	C, 64.48;	H, 5.84;	N, 9.03;	S, 13.77
Found:	C, 64.02;	H, 5.81;	N, 8.82;	S, 13.42.

EXAMPLE 61

5-[(5-Bromo-2-thienyl)methylene]-2-thioxo-4-thiazolidinone, monosodium salt

A suspension of 2.0 g (0.0065 mole) of 5-[(5-bromo-2-thienyl)methylene]-2-thioxo-4-thiazolidinone in 25 ml of ethanol was treated with 3.3 ml of 2.00 N aqueous sodium hydroxide solution. The mixture was warmed on the steam bath until homogeneous and filtered warm. Cooling yielded 1.2 g (56% yield) of the analytically pure salt, mp 288°-dec.

Calcd. for $C_8H_3BrNOS_3Na \cdot 1.5 H_2O$:	C, 27.05;	H, 1.70;	N, 3.94
Found:	C, 27.06;	H, 1.66;	N, 3.70.

EXAMPLE 62

5-[(3-Bromo-4-hydroxy-5-methoxyphenyl)methylene]-2-thioxo-4-thiazolidinone, disodium salt

A suspension of 3.46 g (0.010 mole) of 5-[(3-bromo-4-hydroxy-5-methoxyphenyl)methylene]-2-thioxo-4-thiazolidinone in 50 ml of water was treated with 10.0 ml of 2.00 N aqueous sodium hydroxide solution. The mixture was warmed on the steam bath until nearly homogenous and filtered warm. The cooled filtrate was subjected to vacuum freeze-drying to leave a residue of 3.8 of red solid. The solid was digested briefly on the steam bath with 200 ml of acetone. Filtration gave 3.5 g (88% yield) of the analytically pure salt, mp 310°.

Calcd. for $C_{11}H_6BrNO_3S_2Na_2 \cdot 0.5 H_2O$:	C, 33.09;	H, 1.77;	N, 3.51
Found:	C, 33.03;	H, 1.73;	N, 3.36.

EXAMPLE 63

5-[(4-Hydroxy-3,5-dimethoxyphenyl)methylene]-2,4-thiazolidinedione

Prepared according to the procedure described in Example 1 using 3,5-dimethoxy-4-hydroxybenzaldehyde (4.8 g, 26 mmoles) and 2,4-thiazolidinedione (2.9 g, 25 mmoles). Recrystallization from methanol gave the pure product (1.7 g), mp 248-249° C.


EXAMPLE 64

5-[[3,5-Bis(1,1-dimethylethyl)-4-hydroxyphenyl]methylene]-2,4-thiazolidinedione

A mixture of 3,5-di-t-butyl-4-hydroxybenzaldehyde (6.5 g, 27 mmoles), 2,4-thiazolidinedione (3.0 g, 26 mmoles), sodium acetate (7.6 g, 93 mmoles), and acetic acid (40 ml) is stirred under an inert atmosphere and heated to reflux. After 78 hours the mixture is stirred into water (300 ml), and the precipitate is filtered off, rinsed three times with water, dried, and recrystallized from ethanol. The product is stirred in 1 N NaOH (50 ml) and extracted several times with dichloromethane. The aqueous solution is acidified with acetic acid, stirred for 1-2 hours, and the precipitate is filtered off, rinsed three times with water and dried to afford the

pure product (2.3 g), mp 238-240° C.

The following compounds are prepared by the method described in Example 64:



	Example	R	MP
15	65	H	247-249
	66	3-OMe	197-198
	67	4-OMe	218-220
20	68	3-OMe, 4-OH	225-227

INTERMEDIATE C

2-Thioxo-4-oxazolidinone

30 A solution of 72.9 g (0.75 mole) of potassium thiocyanate and 48.9 g (0.75 mole) of potassium cyanide in 85 ml of water was stirred in an ice bath and treated over 30 minutes with 60 ml (~0.80 mole) of 37% aqueous formaldehyde solution. The rate of addition was adjusted to maintain a reaction temperature of $\leq 10^\circ$. The mixture was stirred for an additional 90 minutes, then treated over 1 hour with 156 ml of concentrated hydrochloric acid (reaction temperature maintained at $\leq 10^\circ$). The mixture was stirred for 16
35 hours as it slowly warmed to room temperature. The inorganic solids were filtered and discarded, and the filtrate was warmed on the steam bath for 90 minutes. The reaction mixture was again filtered, and the filtrate was extracted with ether (5X 200 ml). The combined extracts were dried (anhydrous sodium sulfate) and evaporated to an oil, which slowly crystallized. The solid was washed with hexane to yield 22.7 g (26% yield), mp 106-109°. A sample recrystallized from ethyl acetate/hexane was analytically pure, mp 110-112°
40 (a mp of 113° is given by N. K. Ushenko and T. E. Gorizdra, Ukrain. Khim. Zhur., 16, 545 (1950)).

EXAMPLE 69

5-[(4-Hydroxy-3-methoxyphenyl)methylene]-2-thioxo-4-oxazolidinone

45 A mixture of 3.25 g (0.021 mole) of 4-hydroxy-3-methoxybenzaldehyde, 2.34 g (0.020 mole) of 2-thioxo-4-oxazolidinone, 5.8 g (0.071 mole) of sodium acetate and 15 ml of acetic acid was stirred and heated at reflux under a nitrogen atmosphere for 2 hours. The mixture was then stirred at room temperature for 16 hours, and added to 300 g of ice/water. The precipitated solid was filtered, washed with water, and recrystallized from aqueous methanol/N,N-dimethylformamide to yield 2.4 g (49% yield) of the oxazole product. A sample recrystallized a second time as above was analytically pure, mp 240°-dec. (A mp of
50 237-238° is given by T. E. Gorizdra and S. N. Baranov, Zhur. Obshchei Khim., 26, 3092 (1956)).

Calcd. for $C_{11}H_9NO_4S \cdot 0.25H_2O$:	C, 51.65;	H, 3.74;	N, 5.48;	S, 12.54
Found:	C, 51.54;	H, 3.97;	N, 5.38;	S, 12.17.

Also prepared by the procedure described in Example 69, utilizing the appropriate aldehyde intermediate were:

EXAMPLE 70

5-[(3-Methoxyphenyl)methylene]-2-thioxo-4-oxazolidinone

mp 215° -dec (recrystallized from methanol (N,N-dimethylformamide)).

Calcd. for $C_{11}H_9NO_3S$:	C, 56.15;	H, 3.86;	N, 5.95;	S, 13.63
Found:	C, 56.34;	H, 3.87;	N, 5.95;	S, 13.51.

EXAMPLE 71

5-[(4-Methoxyphenyl)methylene]-2-thioxo-4-oxazolidinone

mp 205-207° (recrystallized from aqueous methanol/N,N-dimethylformamide). (A mp of 192° if given by N. K. Ushenko and T. E. Gorizdra, Ukrain Khim. Zhur., 16, 545 (1950)).

Calcd. for $C_{11}H_9NO_3S$:	C, 56.15;	H, 3.86;	N, 5.95;	S, 13.63
Found:	C, 56.10;	H, 3.88;	N, 6.03;	S, 13.62.

EXAMPLE 72

5-[(3,4-Dimethoxyphenyl)methylene]-2-thioxo-4-oxazolidinone

mp 249° -dec.

Calcd. for $C_{12}H_{11}NO_4S$:	C, 54.32;	H, 4.18;	N, 5.28;	S, 12.09
Found:	C, 53.92;	H, 4.10;	N, 5.22;	S, 11.84.

EXAMPLE 73

5-[(4-Hydroxy-3,5-dimethoxyphenyl)methylene]-2-thioxo-4-oxazolidinone

mp 250° -dec. (recrystallized from aqueous acetonitrile/N,N-dimethylformamide).

Calcd. for $C_{12}H_{11}NO_5S$:	C, 51.24;	H, 3.94;	N, 4.98;	S, 11.40
Found:	C, 51.00;	H, 3.78;	N, 5.30;	S, 11.45.

5

EXAMPLE 74

10 5-[(4-Hydroxy-3,5-dimethylphenyl)methylene]-2-thioxo-4-oxazolidinone

mp 253° -dec.

15

Calcd. for $C_{12}H_{11}NO_3S$:	C, 57.81;	H, 4.45;	N, 5.62;	S, 12.86
Found:	C, 57.84;	H, 4.49;	N, 5.56;	S, 12.54.

20

EXAMPLE 75

5-[(3-Bromo-4-hydroxy-5-methoxyphenyl)methylene]-2-thioxo-4-oxazolidinone

25

mp 260-262°

30

Calcd. for $C_{11}H_8BrNO_4S$:	C, 40.01;	H, 2.44;	N, 4.24;	S, 9.71;	Br, 24.20
Found:	C, 40.18;	H, 2.34;	N, 4.07;	S, 9.61;	Br, 24.02.

35

EXAMPLE 76

2-Thioxo-5-[(2,4,5-trimethoxyphenyl)methylene]-4-oxazolidinone

mp 265° -dec. (recrystallized from aqueous methanol/N,N-dimethylformamide).

40

Calcd. for $C_{13}H_{13}NO_5S$:	C, 52.87;	H, 4.44;	N, 4.74;	S, 10.86
Found:	C, 52.55;	H, 4.30;	N, 4.79;	S, 10.47.

45

EXAMPLE 77

50 2-Thioxo-5-[(3,4,5-trimethoxyphenyl)methylene]-4-oxazolidinone

mp 230° -dec. (recrystallized from aqueous methanol/N,N-dimethylformamide).

55

Calcd. for $C_{13}H_{13}NO_5S$:	C, 52.87;	H, 4.44;	N, 4.74;	S, 10.86
Found:	C, 52.81;	H, 4.32;	N, 4.62;	S, 10.39.

EXAMPLE 78

5 5-[(4-Hydroxy-3,5-dimethoxyphenyl)methylene]-2-thioxo-4-oxazolidinone, disodium salt

A suspension of 3.1 g (0.011 mole) of 5-[(4-hydroxy-3,5-dimethoxyphenyl)methylene]-2-thioxo-4-oxazolidinone in 20 ml of water was treated with a solution of 11.0 ml of 2.00 N aqueous sodium hydroxide solution. The mixture was warmed briefly on the steam bath until nearly homogenous, then filtered warm.
 10 The cooled filtrate was subjected to vacuum freeze-drying to yield 3.5 g (98% yield) of the analytically pure salt product, mp >265°

Calcd. for $C_{12}H_9NO_5Na_2 \cdot 0.6 H_2O$:	C, 42.89;	H, 3.06;	N, 4.17
Found:	C, 42.67;	H, 2.88;	N, 4.00.

Also prepared by the above procedures was:

20 EXAMPLE 79

25 5-[(4-Hydroxy-3-methoxyphenyl)methylene]-2-thioxo-4-oxazolidinone, disodium salt

mp >270°

Calcd. for $C_{11}H_7NO_4Na_2 \cdot 1 H_2O$:	C, 42.18;	H, 2.90;	N, 4.47
Found:	C, 41.84;	H, 2.52;	N, 4.37.

35 EXAMPLE 80

35 5-[(3,5-Dibromo-4-hydroxyphenyl)methylene]-2-thioxo-4-imidazolidinone

Prepared according to the procedure described in Example 1 using 3,5-dibromo-4-hydroxybenzaldehyde (4.5 g, 16 mmoles), 2-thiohydantoin (1.7 g, 15 mmoles), sodium acetate (4.5 g, 55 mmoles), and acetic acid (35 ml). Recrystallization from ethanol gave the pure product (1.4 g), mp 267° C (dec).

45 EXAMPLE 81

50 5-[[4-hydroxy-3,5-bis(1-methylethyl)phenyl]methylene]-2-thioxo-4-imidazolidinone

Prepared according to the procedure described in Example 1 using 4-hydroxy-3,5-bis(1-methylethyl)benzaldehyde (3.0 g, 15 mmoles), 2-thiohydantoin (1.6 g, 14 mmoles), sodium acetate (4.2 g, 51 mmoles), and acetic acid (35 ml), filtered off, dried, and recrystallized from acetonitrile to afford the pure product (1.4 g), mp 225-230° C.

55 EXAMPLE 82

5-[(3-Bromo-4-hydroxyphenyl)methylene]-2-thioxo-4-imidazolidinone

Prepared according to the procedure described in Example 1 using 3-bromo-4-hydroxybenzaldehyde (3.1 g, 15 mmoles), 2-thiohydantoin (1.7 g, 15 mmoles), sodium acetate (4.5 g, 55 mmoles), and acetic acid (35 ml), to afford the pure product (3.4 g), mp 262° C (dec).

EXAMPLE 83

5-[[3,5-Bis(1,1-dimethylethyl)-4-hydroxyphenyl]methylene]-2-thioxo-4-imidazolidinone


Prepared according to the procedure described in Example 1 using 3,4-di-t-butyl-4-hydroxybenzaldehyde (6.2 g, 26 mmoles), 2-thiohydantoin (2.9 g, 25 mmoles), sodium acetate (7.5 g, 91 mmoles), and acetic acid (40 ml). Heating under reflux is maintained for 36 hours. Recrystallization from acetonitrile gave the pure product (4.0 g), mp 279-281° C (dec).

EXAMPLE 84

5-[(2-Methoxyphenyl)methylene]-2-thioxo-4-imidazolidinone

A mixture of 2-methoxybenzaldehyde (2.5 g, 18 mmoles), 2-thiohydantoin (2.0 g, 17 mmoles), sodium acetate (4.0 g, 49 mmoles), acetic anhydride ($\frac{1}{2}$ ml), and acetic acid (15 ml) is stirred under an inert atmosphere and heated to reflux. After 1 hour the mixture is stirred into water (300 ml), and the precipitate is filtered off, rinsed with water three times and dried. The product is triturated in boiling methanol, filtered off, and dried to afford the pure product (3.4 g), mp 234-235° C.

The following examples are prepared from the corresponding benzaldehydes using the procedure described in Example 84.



Example	R ₁	R ₂	R ₃	mp °C	Recryst.
					Solvent
85	OMe	H	H	238-239	MeOH
86	H	OMe	H	266-267	MeOH
87	OMe	OH	H	237-238	MeCN
88	OMe	OH	OMe	268 (dec)	MeOH

EXAMPLE 89

5-[[4-(Acetyloxy)-3-methoxyphenyl]methylene]-2,4-imidazolidinedione

Prepared according to the procedure described in Example 84, using 4-acetoxy-3-methoxybenzaldehyde (4.0 g, 21 mmoles) and hydantoin (2.0 g, 20 mmoles). Recrystallization from methanol/DMF gave the product (1.8 g), mp 275-277° C, retaining 0.15 equivalents of DMF.

EXAMPLE 90

5-[[4-Hydroxy-3-methoxyphenyl]methylene]-2,4-imidazolidinedione

A mixture of 4-hydroxy-3-methoxybenzaldehyde (4.6 g, 30 mmoles), hydantoin (3.0 g, 30 mmoles), β -alanine (1.4 g, 16 mmoles), and acetic acid (40 ml) is stirred under an inert atmosphere and heated to reflux. After 4 hours the mixture is stirred into water (350 ml) and the precipitate is filtered off, rinsed successively with water (3X), ethanol (2X), and ether (2X), and dried. Recrystallization from methanol afforded the pure product (3.0 g), mp 272-273° C.

EXAMPLE 91

5-[[3,5-Dimethoxy-4-hydroxyphenyl]methylene]-2,4-imidazolidinedione

Prepared according to the procedure described in Example 90 using 3,5-dimethoxy-4-hydroxybenzaldehyde (5.6 g, 30 mmoles), hydantoin (3.0 g, 30 mmoles), and β -alanine (1.4 g, 16 mmoles), to afford the pure product (6.0 g), mp 296-297° C.

EXAMPLE 92

5-[[3,5-Bis(1,1-dimethylethyl)-4-hydroxyphenyl]methylene]-2,4-imidazolidinedione

A mixture of 3,5-di-*t*-butyl-4-hydroxybenzaldehyde (7.0 g, 30 mmoles), hydantoin (3.0 g, 30 mmoles), β -alanine (1.4 g, 16 mmoles), and acetic acid (40 ml) is stirred under an inert atmosphere and heated to reflux. After 48 hours the mixture is stirred into water (300 ml), and the precipitate is filtered off, rinsed with water three times, and dried. Recrystallization from acetonitrile gave the pure product (4.8 g), mp 251-252° C.

EXAMPLE 93

5-[[2-Methoxyphenyl]methylene]-3-methyl-2,4-imidazolidinedione

A mixture of 2-methoxybenzaldehyde (3.4 g, 25 mmoles), 3-methyl hydantoin (3.0 g, 21 mmoles), sodium acetate (6.0 g, 73 mmoles), acetic anhydride (3 ml), and acetic acid (20 ml) is stirred under an inert atmosphere and heated to reflux. After 15 hours the mixture is poured into water (200 ml), stirred, and the precipitate filtered off, rinsed three times with water and dried. Recrystallization from acetonitrile gave the pure product (3.1 g), mp 189-190° C.

EXAMPLE 94

5 5-[(3-Methoxyphenyl)methylene]-3-methyl-2,4-imidazolidinedione

Prepared according to the procedure described in Example 93 using 3-methoxybenzaldehyde (3.5 ml, 28 mmoles) and 3-methyl hydantoin (3.0 g, 21 mmoles), to afford the product (3.6 g), mp 203-205 °C.

10

EXAMPLE 95

15 5-[(4-Methoxyphenyl)methylene]-3-methyl-2,4-imidazolidinedione

Prepared according to the procedure described in Example 93 using 4-methoxybenzaldehyde (3.4 g, 25 mmoles) and 3-methyl hydantoin (3.0 g, 21 mmoles). Recrystallization from acetonitrile gave the pure product (2.0 g), mp 221-222 °C.

20

EXAMPLE 96

25

5-[(4-Hydroxy-3-methoxyphenyl)methylene]-3-methyl-2,4-imidazolidinedione

Prepared according to the procedure described in Example 93 using 4-hydroxy-3-methoxybenzaldehyde (3.2 g, 21 mmoles) and 3-methyl hydantoin (3.0 g, 21 mmoles). Recrystallization from ethyl acetate gave the pure product (0.7 g), mp 227-228 °C.

30

EXAMPLE 97

35

5-[(3,5-Dimethoxy-4-hydroxyphenyl)methylene]-3-methyl-2,4-imidazolidinedione

A mixture of 3,5-dimethoxy-4-hydroxybenzaldehyde (3.2 g, 18 mmoles), 3-methyl hydantoin (2.0 g, 18 mmoles), sodium acetate (6.0 g, 73 mmoles), acetic anhydride (3 ml), and acetic acid (25 ml) is stirred under an inert atmosphere and heated to reflux. After 72 hours the mixture is stirred into water (300 ml) and the precipitate is filtered off, rinsed three times with water and dried to give a product (4.0 g) which is a mixture of the desired product and its O-acetate. A mixture of this product (2.1 g), sodium carbonate (3.5 g), DMSO (5 ml), and methanol (50 ml) is stirred and heated to reflux. After 18 hours the mixture is stirred into icewater (400 ml) and acidified with 4N HCl. The precipitate is filtered off, rinsed successively with water (3X), ethanol (2X), and ether (2X), and dried. Recrystallization from methanol/DMF gave the pure product (0.6 g), mp 269-270 °C.

45

50

EXAMPLE 98

55 5-[(3-Methoxyphenyl)methylene]-1-methyl-2,4-imidazolidinedione

A mixture of 2-methoxybenzaldehyde (3.8 g, 28 mmoles), 1-methyl hydantoin (3.0 g 26 mmoles), sodium acetate (8.4 g, 102 mmoles), acetic anhydride (10 ml), and acetic acid (40 ml) is stirred under an inert atmosphere and heated to reflux. After 24 hours the mixture is stirred into water (350 ml), stirred, and

the resulting gum is isolated and washed by decantation, then crystallized from isopropanol. Recrystallization from acetonitrile gave the pure product (1.1 g), mp 195-197 °C.

EXAMPLE 99

5-[(4-Hydroxy-3-methoxyphenyl)methylene]-1-methyl-2,4-imidazolidinedione

A mixture of 4-hydroxy-3-methoxybenzaldehyde (4.6 g, 30 mmoles), 1-methyl hydantoin (3.5 g, 30 mmoles), β -alanine (1.4 g, 16 mmoles), and acetic acid (40 ml) is stirred under an inert atmosphere and heated to reflux. After 8 hours the mixture is stirred into water (300 ml) and the precipitate filtered off, rinsed successively with water (3X), ethanol (2X), and ether (2X), and dried. Recrystallization from acetonitrile gave the pure product (1.1 g), mp 197-198 °C.

EXAMPLE 100

5-[(3,5-Dimethoxy-4-hydroxyphenyl)methylene]-1-methyl-2,4-imidazolidinedione

Prepared according to the procedure described in Example 99 using 3,5-dimethoxy-4-hydroxybenzaldehyde (5.8 g, 31 mmoles) and 1-methyl hydantoin (3.5 g, 30 mmoles). Recrystallization from ethanol gave the pure product (2.8 g), mp 194-196 °C.

EXAMPLE 101

5-[(3,5-Dibromo-4-hydroxyphenyl)methylene]-1-methyl-2,4-imidazolidinedione

Prepared according to the procedure described in Example 99 using 3,5-dibromo-4-hydroxybenzaldehyde (8.4 g, 30 mmoles) and 1-methyl hydantoin (3.5 g, 30 mmoles) to afford the pure product (8.2 g), mp 271-277 °C.

EXAMPLE 102

5-[[4-Hydroxy-3,5-bis(1-methylethyl)phenyl]methylene]-1-methyl-2,4-imidazolidinedione

Prepared according to the procedure described in Example 92 using 3,5-diisopropyl-4-hydroxybenzaldehyde (2.7 g, 13 mmoles) and 1-methyl hydantoin (1.4 g, 12 mmoles). Recrystallization from acetonitrile gave the pure product (0.4 g), mp 197-208 °C.

The following compounds are prepared according to the procedure described in Example 1 using appropriate corresponding starting materials.

EXAMPLE 103

5-[(4-Pyridyl)methylene]-2-thioxo-4-thiazolidinone, mp >300 °.

EXAMPLE 104

5 5-[(2-Thienyl)methylene]-2-thioxo-4-thiazolidinone, mp 227-229°.

EXAMPLE 105

10

5-[(5-Nitro-2-furanyl)methylene]-2-thioxo-4-thiazolidinone, mp 201-202°.

15

EXAMPLE 106

20

5-[(1-Phenyl-1H-indol-2-yl)methylene]-2-thioxo-4-thiazolidinone, mp 282-283°.

EXAMPLE 107

25

5-(3-Bromo-4-hydroxyphenyl)methylene]-2-thioxo-4-oxazolidinone, mp 240° (dec).

Prepared according to the procedure described in Example 69.

30

EXAMPLE 108

35 5-[[3,5-Bis(1,1-dimethylethyl-4-hydroxyphenyl)methylene]-2-thioxo-4-oxazolidinone

A mixture of 14.1 g (0.060 mole) of 3,5-di-tert-butyl-4-hydroxybenzaldehyde, 7.0 g (0.060 mole) of 2-thioxo-4-oxazolidinone, 17.4 g (0.21 mole) of sodium acetate, and 75 ml of acetic acid is stirred and heated at reflux under a nitrogen atmosphere for 20 hours. The cooled reaction mixture is added to 900 g of ice/water and the precipitated product filtered and washed with water. There is obtained 16.2 g (81% yield) of the oxazole product, suitable for further reaction.

A sample of the above crude product is chromatographed over silica gel, using elution with 2.5% ethyl acetate in dichloromethane followed by 25% ethyl acetate. The chromatography product is recrystallized from aqueous acetonitrile to yield the analytically pure oxazole, mp 240° C dec.

45

Calcd. for C ₁₈ H ₂₃ NO ₃ S:	C, 64.83;	H, 6.95;	N, 4.20
Found:	C, 65.00;	H, 6.95;	N, 4.17.

50

EXAMPLE 109

55

5-[[3,5-Bis(1,1-dimethylethyl-4-hydroxyphenyl)methylene]-2,4-oxazolidinedione

A solution of 10.0 g (0.030 mole) of 5-[[3,5-bis(1,1-dimethylethyl-4-hydroxyphenyl)methylene]-2-thioxo-4-

oxazolidinone in 150 ml of tetrahydrofuran is cooled in ice and treated with 4.2 ml (3.0 g, 0.030 mole) of triethylamine. The mixture is stirred with ice cooling for one hour, then treated with 10.0 ml (22.8 g, 0.16 mole) of iodomethane. The ice bath is removed and the mixture is stirred for an additional 24 hours. The reaction mixture is filtered and the filter cake is washed several times with fresh tetrahydrofuran. The combined filtrates are evaporated and the residue is chromatographed (silica gel, 2% methanol in dichloromethane elution) to yield 4.4 g (42% yield) of the purified 2-methylthio- intermediate. This material is hydrolyzed without further purification.

A solution of 3.0 g (0.086 mole) of the above 2-methylthio-intermediate in 90 ml of ethanol is diluted with 30 ml of water and treated dropwise over 10 minutes with 5.0 ml of concentrated hydrochloric acid. The reaction mixture is stirred for 24 hours and the precipitated product is filtered and washed with hexane. The crude yield is 1.44 g (53%). A sample recrystallized from ethyl acetate/hexane yielded the analytically pure oxazolidinedione, mp 239° C dec.

Calcd. for $C_{18}H_{23}NO_4$:	C, 68.12;	H, 7.31;	N, 4.41
Found:	C, 68.22;	H, 7.29;	N, 4.05.

Also prepared by the above procedure, utilizing the appropriate 2-thioxo-4-oxazolidinone intermediate are:

EXAMPLE 110

5-[(4-Hydroxy-3,5-dimethoxyphenyl)methylene]-2,4-oxazolidinedione

mp 268° dec. (recrystallized from acetonitrile/DMF/water).

Calcd. for $C_{12}H_{11}NO_6$:	C, 54.34;	H, 4.18;	N, 5.28
Found:	C, 54.00;	H, 4.17;	N, 5.15.

EXAMPLE 111

5-[(4-Hydroxy-3,5-dimethylphenyl)methylene]-2,4-oxazolidinedione

mp 275° C dec. (recrystallized from acetonitrile/DMF/water).

Calcd. for $C_{12}H_{11}NO_4 \cdot 0.5H_2O$:	C, 59.50;	H, 4.99;	N, 5.78
Found:	C, 59.31;	H, 5.02;	N, 5.72.

The usefulness of the compounds of the present invention as inhibitors of histamine release is demonstrated by the following assay. The assay is essentially as generally accepted among the ordinarily skilled artisans to show activity having usefulness to treat the diseases or conditions as set out in the present invention. A description of the procedure follows.

HISTAMINE RELEASE FROM HUMAN BASOPHILS

(hereinafter HHB)

The HHB assay quantitates active histamine release, and its inhibition by drugs, from basophils of human blood. Thus, the assay provides evaluation of the compounds of formula I for treating the conditions or diseases as is the present invention. As described herein the assay includes modifications of the method described by R. P. Siraganian in "An Automated Continuous-Flow System for the Extraction and Fluorometric Analysis of Histamine", Anal. Biochem., 57, 383-394 (1974).

EXAMPLE 112

5-[[3,5-Bis(1,1-dimethylethyl)-4-hydroxyphenyl]methylene]-2-thioxo-4-oxazolidinone, choline salt

A solution of 2.45 g (0.0069 mole) of 46.6% aqueous choline bicarbonate in 50 ml of methanol is stirred and treated over 5 minutes with 2.31 g (0.0069 mole) of 5[[3,5-bis(1,1-dimethylethyl)-4-hydroxyphenyl]-methylene]-2-thioxo-4-oxazolidinone. The reaction mixture is warmed for a few minutes to reflux on the steam bath, and filtered hot. The cooled filtrate is evaporated, and the residue recrystallized from acetone/tert-butyl methyl ether to yield 2.1 g (70% yield) of the analytically pure choline salt, mp 167° dec.

Calc. for $C_{23}H_{36}N_2O_4S$:	C, 63.27;	H, 8.31;	N, 6.42
Found:	C, 63.22;	H, 8.21;	N, 6.28

EXAMPLE 113

5-[[3,5-bis(1,1-dimethylethyl)-4-hydroxyphenyl]methylene]-2,4-thiazolidinedione, choline salt

A solution of 46.6% aqueous choline bicarbonate (10.55 g, 29.8 mmoles) is added dropwise under nitrogen to a stirred suspension of 5-[[3,5-bis(1,1-dimethylethyl)-4-hydroxyphenyl]-methylene]-2,4-thiazolidinedione (10.0 g, 30.0 mmoles) in ethanol (100 ml). The solution is gradually warmed to reflux for one hour, and is then cooled and stripped of solvent by rotary evaporator. Two 50 ml portions of ethanol are successively mixed with and stripped from the residue which is then thoroughly dried under vacuum. The residue is triturated in diethyl ether, filtered off, washed twice with ether, and dried under vacuum at 80° C overnight to afford the crystalline choline salt (12.8 g), mp 215° C dec.

Calc. for $C_{18}H_{22}NO_3S \cdot C_5H_{14}NO$:	C, 63.27;	H, 8.31;	N, 6.42
Found:	C, 63.12;	H, 8.28;	N, 6.17

METHODS

Preparation of Leukocytes

Blood is drawn from allergic donors (chosen on the basis of adequate histamine release induced by a challenge), using standard venipuncture methods, into Vacutainers with EDTA in water as anticoagulant. The blood samples are placed briefly on a rotary mixer. The blood is mixed with Hespan (hydroxy ethyl starch, 0.5 ml per 1.0 ml of blood), the tube inverted several times to mix and then left undisturbed at room temperature until a sharp separation is observed between the settled red cells and the leukocyte and platelet-rich plasma. This usually occurs within 35-45 minutes.

The plasma fraction is removed and centrifuged for 12 minutes at 4° C at 1050 RPM (100 Xg). The platelets remain in the plasma and are discarded. The pelleted leukocytes are shaken gently to disrupt the cell button and washed twice with HA buffer containing 0.005 M EDTA and resuspended in HACM. buffer to

approximately one-quarter the original blood volume. A sample is prepared for Hematology, where a total white blood cell and platelet count is done using a Coulter Counter.

5 Protocol Design

Aliquots (0.1 ml) of cells are added to triplicate assay tubes containing 0.4 ml of either 6% perchloric acid (for total histamine content), vehicle control (for spontaneous release), or drug. The tubes are incubated at room temperature for 8 minutes, and then placed in a 37 °C water bath for 2 more minutes. Buffer or challenge agents (at 37 °C) are added to the tubes and they are incubated for an additional 45 minutes at 37 °C in a shaking water bath. The tubes are then spun at 2000 RPM (1200 g) for 3 minutes to pellet the cells and the supernatants are removed and assayed for histamine by the fluorometric method.

15 Drug Preparation

A 300 µM stock solution of each test compound is prepared in distilled water, using 0.5 ml DMSO/100 ml and/or 0.2 ml of 1N NaOH or HCl and/or heat to aid in dissolution. Five ml of the stock solution is diluted (1:2) with 5 ml of two times concentrated HACM buffer to yield the stock working concentration of 150 µM. When added to the cells and stimulus, a final test concentration of 100 µM drug results. Further dilutions are made with HACM buffer for 33, 10, 3.3, 1.0 µM, etc.

Challenge Agent Preparation

Short ragweed and house dust extracts (Greer Laboratories, Inc.) are supplied as aqueous extracts in stock concentrations of 40,000 and 10,000 protein nitrogen units per milliliter (PNU/ml), respectively. Aqueous solutions of anti-IgE antisera (rabbit-raised antibody) are purchased from Dako via Accurate Chemicals. The aqueous solutions of ragweed, house dust, and anti-IgE are diluted 1:2 with two times concentrated HACM and then further diluted with HACM to yield final stock concentrations of 6000 PNU/ml for ragweed and house dust and 1:50 dilution for the anti-IgE antisera. Further dilutions for working solutions are made in HACM buffer. All stock and working solutions are stored at 4 °C. Working solutions comprise 1/6 of the final volume in the cell reaction, therefore, working solutions of challenge agents are made up six times the required final concentration.


In each experiment, cells are challenged according to the previously determined sensitivity of that donor to the particular challenge agent. Short ragweed and house dust concentrations are expressed in PNU/ml, and anti-IgE antisera is expressed as dilutions, e.g., IE-5 (1:100,000), 3E-5 (1:30,000), and IE-4 (1:10,000).

40 Calculation and Interpretation of Results

The total histamine concentration in the "total" (acid-treated) samples must be 15 ng/ml to be acceptable. Spontaneous release of histamine from the cells should not exceed 15% of the total histamine, and is frequently < 5%. The maximum percentage histamine released varied with the donor. The net amount released by the challenge agent must exceed 25% of the total cellular histamine to confidently assess inhibition by test compounds. Spontaneous histamine release is subtracted from both "totals" and challenged cells to calculate net percent release. Percent inhibition is shown in the Table and is calculated using the following formula:

$$1 - \left(\frac{\text{Mean net \% release treated samples}}{\text{Mean net \% release for challenged control}} \right) \times 100 = \% \text{ inhibition}$$

TABLE



Ex	R ₁	R ₂	R ₃	R ₄	R ₅	% Inhibition of ^{HHB} Histamine Release at 33 μ M
1	MeO	MeO	H	H	H	*
2	MeO	H	OMe	H	H	*
3	MeO	H	H	OMe	H	86
4	MeO	OMe	OMe	H	H	*
5	MeO	H	OMe	OMe	H	73
6	H	OMe	OMe	OMe	H	92
7	Cl	H	Cl	H	H	81
8	H	Cl	Cl	H	H	88
9	H	OMe	OMe	H	H	84
10	H	OMe	OH	OMe	H	90
11	H	F	OMe	H	H	94
12	H	OH	OH	H	H	41
13	H	Br	OH	H	H	61
14	H	Br	OH	Br	H	*
15	H	Me	OH	Me	H	84
16	H	OMe	OH	Br	H	51
17	H	H	H	H	H	39
18	MeO	H	H	H	H	*
19	H	MeO	H	H	H	100
20	H	H	MeO	H	H	90


						<u>HMB</u>
						% Inhibition of Histamine Release
						at 33 μ M
Ex	R ₁	R ₂	R ₃	R ₄	R ₅	
5	21	H	OH	H	H	75
	22	H	H	OH	H	95
	23	H	OEt	H	H	*
10	24	H	OCHMe ₂	H	H	*
	25	H	OCH ₂ Ph	H	H	*
	26	H	Me	H	H	58
	27	H	CF ₃	H	H	58
15	28	H	Cl	H	H	*
	29	H	Br	H	H	*
	30	H	NO ₂	H	H	*
20	31	H	H	Cl	H	95
	32	H	H	Br	H	61
	33	H	H	OPh	H	64
	34	H	H	NO ₂	H	*
25	35	H	H	SMe	H	*
	36	H	MeO	OH	H	79
	37	H	MeO	H	MeO	80
30	38	H	H	Me	H	30
	39	H	H	Ph	H	28
	40	Cl	Cl	H	H	60
	41	Cl	H	H	H	*
35	43	H	F	H	H	*
	44	OH	H	H	H	*

40

45

50

55



5

10


Ex	<u>R₁</u>	<u>R₂</u>	<u>R₃</u>	<u>R₄</u>	<u>R₅</u>	<u>HHB</u> % Inhibition of Histamine Release at 33 μ M
15	45	H	OH	OMe	H	59
	47	H	OH	OMe	OMe	79
	48	OMe	H	H	OMe	*
20	49	OH	H	H	OMe	98
	50	H	OH	H	OMe	96
	51	H	OMe	OH	tBu	34
	52	OH	H	OMe	OMe	79
25	53	H	Me ₂ NCH ₂	OH	MeO	*

30


	<u>HHB</u>
	% Inhibition of Histamine Release
<u>Ex</u>	<u>at 33 μM</u>
54	85
35	88
56	26
57	66
58	24
40	72
59	95
103	87
104	88
45	84
105	
106	

50

55




Ex	<u>R₁</u>	<u>R₂</u>	<u>R₃</u>	<u>R₄</u>	<u>R₅</u>	<u>HHB</u> % Inhibition of Histamine Release at 33 μ M
63	H	MeO	OH	MeO	H	60
64	H	tBu	OH	tBu	H	49
65	H	H	H	H	H	*
66	H	OMe	H	H	H	92
67	H	H	OMe	H	H	90
68	H	OMe	OH	H	H	81




Ex	<u>R₁</u>	<u>R₂</u>	<u>R₃</u>	<u>R₄</u>	<u>R₅</u>	<u>HHB</u> % Inhibition of Histamine Release at 33 μ M
69	H	MeO	OH	H	H	77
70	H	MeO	H	H	H	58
71	H	H	MeO	H	H	83
72	H	MeO	MeO	H	H	43
73	H	MeO	OH	MeO	H	89

							HHB
							% Inhibition of Histamine Release
							at 33 μ M
<u>Ex</u>	<u>R₁</u>	<u>R₂</u>	<u>R₃</u>	<u>R₄</u>	<u>R₅</u>		
5	74	H	Me	OH	Me	H	88
	75	H	Br	OH	MeO	H	*
	76	MeO	H	MeO	MeO	H	*
10	77	H	MeO	MeO	MeO	H	76
	107	H	Br	OH	H	H	66



						<u>HHB</u> % Inhibition of Histamine Release at 33 μ M	
	<u>Ex</u>	<u>R₁</u>	<u>R₂</u>	<u>R₃</u>	<u>R₄</u>	<u>R₅</u>	
30	80	H	Br	OH	Br	H	85
	82	H	Br	OH	H	H	78
35	83	H	tBu	OH	tBu	H	30
	84	OMe	H	H	H	H	*
	85	H	OMe	H	H	H	28
40	86	H	H	OMe	H	H	*
	87	H	OMe	OH	H	H	*
	88	H	OMe	OH	OMe	H	68




5

10

Ex	<u>R₁</u>	<u>R₂</u>	<u>R₃</u>	<u>R₄</u>	<u>R₅</u>	<u>HMB</u> % Inhibition of Histamine Release at 33 μ M
89	H	MeO	AcO	H	H	27
90	H	MeO	OH	H	H	*
91	H	MeO	OH	MeO	H	*
92	H	tBu	OH	tBu	H	11

20

25



30


35

Ex	<u>R₁</u>	<u>R₂</u>	<u>R₃</u>	<u>R₄</u>	<u>R₅</u>	<u>HMB</u> % Inhibition of Histamine Release at 33 μ M
93	MeO	H	H	H	H	*
94	H	MeO	H	H	H	*
95	H	H	MeO	H	H	*
96	H	MeO	OH	H	H	58
97	H	MeO	OH	MeO	H	51

45

50

55



5

10

Ex	R ₁	R ₂	R ₃	R ₄	R ₅	<u>HHB</u> % Inhibition of Histamine Release at 33 μ M
98	H	MeO	H	H	H	*
99	H	MeO	OH	H	H	92
100	H	MeO	OH	MeO	H	68
101	H	Br	OH	Br	H	96

15

20

* Indicates not active at 33 μ M.

E Enhanced release at this concentration.

25

MeO is methoxy.

OCHMe₂ is isopropoxy.

OCH₂Ph is benzyloxy.

30

tBu is tertiarybutyl.

Me₂NCH₂ is dimethylamine.

Activity is measured as % inhibition of histamine release from human basophils challenged with anti-IgE at a 33 μ M concentration of drug.

The IC₅₀ is calculated to give the μ M necessary to provide 50% inhibition.

The usefulness of the compounds of the present invention as inhibitors of the 5-lipoxygenase enzyme, cyclooxygenase, or in treating related diseases or conditions may be demonstrated by their effectiveness in various standard test procedures. A description of each procedure follows.

40

ARBL/ARBC Whole Cell 5-Lipoxygenase and Cyclooxygenase Assays

45

Materials

The rat basophilic leukemia cell line (RBL-1) was obtained from the American Type Culture Collection (Rockville, MD).

Radioimmunoassay (RIA) kits of LTB₄ and PGF_{2 α} were obtained from Amersham (Arlington Heights, IL) and Seragen (Boston, MA), respectively.

All tissue culture media were obtained from GIBCO (Grand Island, NY).

Method

55

RBL-1 cells are grown in suspension culture in Eagle's minimum essential medium supplemented with 12% fetal bovine serum at 37° C in an incubator supplied with air-5% carbon dioxide. Cells are harvested

by centrifugation. They are washed with cold phosphate buffered saline pH 7.4 (PBS; NaCl, 7.1 g; Na₂HPO₄, 1.15 g; KH₂PO₄, 0.2 g; and KCl, 0.2 g/l). Cells are finally suspended in PBS containing 1.0 mM calcium at a density of 2×10^6 cells/ml. Cells are incubated with and without test agent (in DMSO) (1% DMSO is without effect on arachidonic acid metabolism) for ten minutes at room temperature. Calcium ionophore A23187 (5 μ M) is added and cells are incubated for seven minutes at 37° C. The reaction is stopped by chilling the tubes on ice for ten minutes. Cells are separated by centrifugation and the supernatant is stored at -20°. Aliquots (100 μ l) are analyzed for LTB₄ and PGF_{2 α} using radioimmunoassay kits as provided by the supplier.

Table 1 contains biochemical data obtained from this whole cell assay as IC₅₀s which are calculated as the amount of test compound causing 50% inhibition of LTB₄ or PGF_{2 α} formation.

Carrageenan-Induced Rat Foot Paw Edema-2

(CEF-2) Assay: Protocol

Carrageenan solution (1% w/v) is prepared by dissolving 100 mg carrageenan (Marine Colloidal Div., Springfield, NJ) in 10 ml of sterile saline (0.9%) solution (Travenol). The solution is vortexed for 30 to 45 minutes. Animals are dosed with compound one hour before carrageenan challenge. Foot paw edema is induced by injecting 0.10 ml of the 1 % carrageenan subcutaneously into the plantar portion of the right hind paw of each rat under light anesthesia. Initial foot paw volume is measured immediately following carrageenan challenge using mercury plethysmography (Buxco Electronics). Edema is measured five hours after carrageenan. The difference between the five-hour and the initial paw volume is expressed as delta edema. The delta edema for each test group of animals is used to calculate the percent inhibition of edema achieved by the compound at the test dose compared with the vehicle control group. The ID₄₀ (the dose at which swelling is inhibited by 40%) is calculated by probit analysis for the dose at which 40 percent inhibition occurs.

TABLE II

Example No.	ARBL ⁽¹⁾	ARBC ⁽²⁾	CFE ⁽³⁾	
64	100	95	31.8	(1)
			39.8	(3)
			53.7	(10)
			58.8	(30)
108	92	86	44.1	(1)
			41.1	(3)
			32.1	(10)
			40.4	(30)
109	95	89	18.5	(1)
			23.5	(3)
			36.8	(10)
			39.4	(30)
110	67	N ⁽⁴⁾		
32	93	59		

(1) Percent inhibition of cellular 5-lipoxygenase at 16 μ M.

(2) Percent inhibition of cellular 5-cyclooxygenase at 16 μ M.

(3) Percent inhibition of carrageenan footpad edema (CFE) test at various doses (mg/kg) of test drug.

(4) N inactive at screening concentration.

Accordingly, the present invention also includes a pharmaceutical composition for treating one of the

above diseases or conditions comprising an antidisease or anticondition effective amount of a compound of the formula I as defined above together with a pharmaceutically acceptable carrier.

The present invention further includes a method for treating one of the above named diseases or conditions in mammals, including man, suffering therefrom comprising administering to such mammals
 5 either orally or parenterally, preferably oral, a corresponding pharmaceutical composition containing a compound of formula I as defined above in appropriate unit dosage form.

For preparing pharmaceutical compositions from the compounds described by this invention, inert, pharmaceutically acceptable carriers can be either solid or liquid. Solid form preparations include powders, tablets, dispersible granules, capsules, cachets, and suppositories. A solid carrier can be one or more
 10 substances which may also act as diluents, flavoring agents, solubilizers, lubricants, suspending agents, binders or tablet disintegrating agents; it can also be encapsulating material. In powders, the carrier is a finely divided active compound. In the tablet the active compound is mixed with carrier having the necessary binding properties in suitable proportions and compacted in the shape and size desired. The powders and tablets preferably contain from 5 or 10 to about 70 percent of the active ingredient. Suitable
 15 solid carriers are magnesium carbonate, magnesium stearate, talc, sugar, lactose, pectin, dextrin, starch, gelatin, tragacanth, methylcellulose, sodium carboxymethylcellulose, a low melting wax, cocoa butter, and the like. The term "preparation" is intended to include the formulation of the active compound with encapsulating material as carrier providing a capsule in which the active component (with or without other carriers) is surrounded by carrier, which is thus in association with it. Similarly, cachets are included.
 20 Tablets, powders, cachets, and capsules can be used as solid dosage forms suitable for oral administration.

For preparing suppositories, a low melting wax such as a mixture of fatty acid glycerides or cocoa butter is first melted, and the active ingredient is dispersed homogeneously therein as by stirring. The molten homogeneous mixture is then poured into convenient sized molds, allowed to cool and thereby to solidify.

25 Liquid form preparations include solutions, suspensions, and emulsions. As an example may be mentioned water or water-propylene glycol solution for parenteral injection. Liquid preparations can also be formulated in solution in aqueous polyethylene glycol solution. Aqueous solutions suitable for oral use can be prepared by dissolving the active component in water and adding suitable colorants, flavors, stabilizing and thickening agents as desired. Aqueous suspensions suitable for oral use can be made by dispersing
 30 the finely divided active component in water with viscous material, i.e., natural or synthetic gums, resins, methylcellulose, sodium carboxymethylcellulose, and other well-known suspending agents.

Also included are solid form preparations which are intended to be converted, shortly before use, to liquid form preparations for either oral or parenteral administration. Such liquid forms include solutions, suspensions, and emulsions. These particular solid form preparations are most conveniently provided in unit
 35 dose form and as such are used to provide a single liquid dosage unit. Alternatively, sufficient solid may be provided so that after conversion to liquid form, multiple individual liquid doses may be obtained by measuring predetermined volumes of the liquid form preparation as with a syringe, teaspoon, or other volumetric container. When multiple liquid doses are so prepared, it is preferred to maintain the unused portion of said liquid doses at low temperature (i.e., under refrigeration) in order to retard possible
 40 decomposition. The solid form preparations intended to be converted to liquid form may contain, in addition to the active material, flavorants, colorants, stabilizers, buffers, artificial and natural sweeteners, dispersants, thickeners, solubilizing agents, and the like. The liquid utilized for preparing the liquid form preparation may be water, isotonic water, ethanol, glycerine, propylene glycol, and the like as well as mixtures thereof. Naturally, the liquid utilized will be chosen with regard to the route of administration, for example, liquid
 45 preparations containing large amounts of ethanol are not suitable for parenteral use.

Preferably, the pharmaceutical preparation is in unit dosage form. In such form, the preparation is subdivided into unit doses containing appropriate quantities of the active component. The unit dosage form can be a packaged preparation, the package containing discrete quantities of preparation, for example, packeted tablets, capsules, and powders in vials or ampoules. The unit dosage form can also be a capsule,
 50 cachet, or tablet itself or it can be the appropriate number of any of these in packaged form.

The quantity of active compound in a unit dose of preparation may be varied or adjusted from 1 mg to 500 mg preferably 1 to 50 mg according to the particular application and the potency of the active ingredient. The compositions can, if desired, also contain other compatible therapeutic agents.

In therapeutic use as described above, the dosages may be varied depending upon the requirements of
 55 the patient, the severity of the condition being treated, and the compound being employed. Determination of the proper dosage for a particular situation is within the skill of the art. Generally, treatment is initiated with the smaller dosages which are less than the optimum dose of the compound. Thereafter the dosage is increased by small increments until the optimum effect under the circumstances is reached. For conve-

nience, the total daily dosage may be divided and administered in portions during the day if desired.

In addition to the compounds of formula I, the pharmaceutical compositions can also contain other active ingredients, such as cyclooxygenase inhibitors, nonsteroidal antiinflammatory drugs (NSAIDs), peripheral analgesic agents such as zomepirac, diflunisal, and the like. The weight ratio of the compound of the formula I to the second active ingredient may be varied and will depend upon the effective dose of each ingredient. Generally, an effective dose of each will be used. Thus, for example, when a compound of the formula I is combined with a NSAID, the weight ratio of the compound of the formula I to the NSAID will generally range from about 1000:1 to about 1:1000, preferably about 200:1 to about 1:200. Combinations of a compound of the formula I and other active ingredients will generally also be within the aforementioned range, but in each case, an effective dose of each active ingredient should be used.

Combinations of a compound of the formula I and other active ingredients will generally be in the aforementioned ratios.

NSAIDs can be characterized into five groups:

- (1) the propionic acid derivatives;
- (2) the acetic acid derivatives;
- (3) the fenamic acid derivatives;
- (4) the biphenylcarboxylic acid derivatives; and
- (5) the oxicams


or a pharmaceutically acceptable salt thereof.

The propionic acid derivatives which may be used comprise: ibuprofen, ibuprofen aluminum, indoprofen, ketoprofen, naproxen, benoxaprofen, flurbiprofen, fenoprofen, fenbufen, piroprofen, carprofen, oxaprozin, pranoprofen, miroprofen, tiaprofen, alminoprofen, tiaprofen, fluprofen, and buclocic acid. Structurally related propionic acid derivatives having similar analgesic and antiinflammatory properties are also intended to be included in this group.

Thus, "propionic acid derivatives" as defined herein are nonnarcotic analgesics/nonsteroidal antiinflammatory drugs having a free $-\text{CH}(\text{CH}_3)\text{COOH}$ or $-\text{CH}_2\text{CH}_2\text{COOH}$ group (which optionally can be in the form of a pharmaceutically acceptable salt group, e.g., $-\text{CH}(\text{CH}_3)\text{COO}^-\text{Na}^+$ or $-\text{CH}_2\text{CH}_2\text{COO}^-\text{Na}^+$), typically attached directly or via a carbonyl function to a ring system, preferably to an aromatic ring system.

The acetic acid derivatives which may be used comprise: indomethacin, which is a preferred NSAID, sulindac, tolmetin, zomepirac, diclofenac, fenclofenac, alclofenac, ibufenac, isoxepac, furofenac, tiopinac, zidometacin, acemetacin, fentiazac, clidanac, oxpinac, and fenclozic acid. Structurally related acetic acid derivatives having similar analgesic and antiinflammatory properties are also intended to be encompassed by this group.


Thus, "fenamic acid derivatives" as defined herein are nonnarcotic analgesics/nonsteroidal antiinflammatory drugs which contain the basic structure:



which can bear a variety of substituents and in which the free $-\text{COOH}$ group can be in the form of a pharmaceutically acceptable salt group, e.g., $-\text{COO}^-\text{Na}^+$.

The biphenylcarboxylic acid derivatives which can be used comprise: diflunisal and flufenisal. Structurally related biphenylcarboxylic acid derivatives having similar analgesic and antiinflammatory properties are also intended to be encompassed by this group.

Thus, "biphenylcarboxylic acid derivatives" as defined herein are nonnarcotic analgesics/nonsteroidal antiinflammatory drugs which contain the basic structure:




which can bear a variety of substituents and in which the free $-\text{COOH}$ group can be in the form of a pharmaceutically acceptable salt group, e.g., $-\text{COO}^-\text{Na}^+$.

The oxicams which can be used in the present invention comprise: piroxicam, sudoxicam, isoxicam,

and 4-hydroxyl-1,2-benzothiazine 1,1-dioxide 4-(N-phenyl)-carboxamide. Structurally related oxicams having similar analgesic and antiinflammatory properties are also intended to be encompassed by this group.

Thus "oxicams" as defined herein are nonnarcotic analgesics/nonsteroidal antiinflammatory drugs which have the general formula:



wherein R is an aryl or heteroaryl ring system.

The following NSAIDs may also be used: acetaminophen, alminoprofen, amfenac sodium, aminoprofen, anitrazafen, antrafenine, auranofin, bendazac lysinate, benzydamine, beprozoin, broperamole, bufezolac, carprofen, cinmetacin, ciproquazone, clidanac, cloximate, dazidamine, deboxamet, delmetacin, detomidine, dexindoprofen, diacerein, di-fisalamine, difenpyramide, ermofazone, enfenamic acid, enolicam, eprizole, etersalate, etodolac, etofenamate, fanetizole mesylate, fenclofenac, fenclorac, fendosal, fenflumizole, fentiazac, feprazone, floctafenine, flunixin, flunoxaprofen, fluproquazone, fopirtoline, fosfosol, furciprofen, furofenac, glucametacin, guimesal, ibuprofen, isofezolac, isonixim, isoprofen, isoxepac, isoxicam, lefetamine HCl, leflunomide, lofenizole, lonazolac calcium, lotifazole, loxoprofen, lysin, clonixinate, meclofenamate sodium, meseclozone, microprofen, nabumetone, nictindole, nimesulide, orpanoxin, oxametacin, oxapadol, oxaprozin, perisoxal citrate, pimeprofen, pimetacin, piroxifen, pirazolac, pirfenidone, pirprofen, pranoprofen, proglumetacin maleate, proquazone, pyridoxiprofen, sudoxicam, suprofen, talmetacin, talniflumate, tenoxicam, thiazolinobutazone, thielavin B, tiaprofenic acid, tiaramide HCl, tifenizole, timogadine, tioprofen, tolfenamic acid, tolpadol, tryptamid, ufenamate, and zidometacin.


Finally, NSAIDs which may also be used include the salicylates, specifically aspirin, and the phenylbutazones, and pharmaceutically acceptable salts thereof.

Pharmaceutical composition comprising the formula I compounds may also contain as the second active ingredient, antihistaminic agents such as benadryl, dramamine, histadyl, phenergan, and the like. Alternatively, they may include prostaglandin antagonists such as those disclosed in European Patent Application 11,067 or thromboxane antagonists such as those disclosed in U.S. 4,237,160. They may also contain histidine decarboxylase inhibitors such as α -fluoromethylhistidine, described in U.S. 4,325,961. The compounds of formula I may also be advantageously combined with an H_1 or H_2 -receptor antagonist, such as for instance cimetidine, ranitidine, terfenadine, famotidine, temelastine, acrivastine, loratidine, cetirizine, tazifylline, azelastine, aminothiadiazoles disclosed in EP 81102976.8 and like compounds, such as those disclosed in U.S. Patent Nos. 4,283,408; 4,362,736; 4,394,508, and European Patent Application No. 40,696. The pharmaceutical compositions may also contain a K^+/H^+ ATPase inhibitor such as omeprazole, disclosed in U.S. Patent 4,255,431, and the like. Each of the references referred to in this paragraph is hereby incorporated herein by reference.

Thus, "acetic acid derivatives" as defined herein are nonnarcotic analgesics/nonsteroidal antiinflammatory drugs having a free $-CH_2COOH$ group (which optionally can be in the form of a pharmaceutically acceptable salt group, e.g., $-CH_2COO^-Na^+$), typically attached directly to a ring system, preferably to an aromatic or heteroaromatic ring system.

Claims

1. A pharmaceutical composition comprising a compound of the formula (I)



and pharmaceutically acceptable salts thereof;

wherein Ar is (i) phenyl;

(ii) phenyl substituted by from one to three of lower alkyl, lower alkoxy, hydroxy, halogen, trifluoromethyl, NO₂, mercapto, or lower alkylthio, NR₁₀R₁₁ wherein R₁₀ and R₁₁ are independently hydrogen, or lower alkyl;

(iii) naphthyl;

(iv) benzofuranyl;

(v) benzothiophenyl;

(vi) thienyl;

(vii) indolyl

(viii) furanyl

(ix) pyridyl


Y and Y₁ is oxygen or sulfur;

X is sulfur, oxygen, NH or NCH₃; and

X₁ is NH or NCH₃;


with the proviso that Ar is not 3,5-di(isopropyl)-4-hydroxyphenyl and further if X₁ is NH, X is S, and Y is S then Ar is not 3,5-di(t-butyl)-4-hydroxyphenyl and a pharmaceutically acceptable carrier.

2. A composition of Claim 1 wherein the compound is of the formula (II)




wherein X and Y are as defined above and having of from one to three of R₁, R₂, R₃, R₄ and R₅ selected from the group consisting of lower alkyl, lower alkoxy, OH, halogen, CF₃, NR₁₀R₁₁ wherein R₁₀ and R₁₁ are as defined above, NO₂ mercapto, or lower alkylthio.

3. A composition of Claim 2 wherein the compound is of the formula (III)




wherein R₁, R₂, R₃, R₄ and R₅ are as defined above.

4. A composition of Claim 2 wherein the compound is of the formula (IV)




5. A composition of Claim 2 wherein the compound is of the formula (V)




wherein R_1 , R_2 , R_3 , R_4 and R_5 are as defined above.

6. A composition of Claim 2 wherein the compound is of the formula (VI)




wherein R_1 , R_2 , R_3 , R_4 and R_5 are as defined above.

7. A composition of Claim 2 wherein the compound is of the formula (VII)




wherein R_1 , R_2 , R_3 , R_4 and R_5 are as defined above.

8. A composition of Claim 1 wherein the compound is of the formula (VIII)



wherein R_1 , R_2 , R_3 , R_4 and R_5 are as defined above.

9. A composition of Claim 2 wherein the compound is of the formula (VIII A)



wherein R_1 , R_2 , R_3 , R_4 and R_5 are as defined above.


10. A composition of Claim 3 wherein the compound is a member of the group consisting of
 5-[(2,5-dimethoxyphenyl)methylene]-2-thioxo-4-thiazolidinone;
 5-[(2,4,5-trimethoxyphenyl)methylene]-2-thioxo-4-thiazolidinone;
 the monosodium salt of 5-[2,4,5-trimethoxyphenyl)methylene]-2-thioxo-4-thiazolidinone;

- 5-[(3,4,5-trimethoxyphenyl)methylene]-2-thioxo-4-thiazolidinone;
 5-[(3,4-dichlorophenyl)methylene]-2-thioxo-4-thiazolidinone;
 the monosodium salt of 5-[(3,4,5-trimethoxyphenyl)methylene]-2-thioxo-4-thiazolidinone;
 5-[(3,4-dimethoxyphenyl)methylene]-2-thioxo-4-thiazolidinone;
 5 the monosodium salt of 5-[(3,4-dichlorophenyl)methylene]-2-thioxo-4-thiazolidinone;
 5-[(3,5-dimethoxy-4-hydroxyphenyl)methylene]-2-thioxo-4-thiazolidinone;
 the disodium salt of 5-[(3,4,5-trimethoxyphenyl)methylene]-2-thioxo-4-thiazolidinone;
 5-[(3,5-dimethyl-4-hydroxyphenyl)methylene]-2-thioxo-4-thiazolidinone;
 the disodium salt of 5-[(3,4-dimethoxyphenyl)methylene]-2-thioxo-4-thiazolidinone;
 10 5-[(3-bromo-4-hydroxy-5-methoxyphenyl)methylene]-2-thioxo-4-thiazolidinone;
 the disodium salt of 5-[(3,4-dichlorophenyl)methylene]-2-thioxo-4-thiazolidinone;
 5-[(4-methoxyphenyl)methylene]-2-thioxo-4-thiazolidinone;
 5-[(3-hydroxy-4-methoxyphenyl)methylene]-2-thioxo-4-thiazolidinone;
 5-[(3,5-dimethoxyphenyl)methylene]-2-thioxo-4-thiazolidinone.
11. A composition of Claim 3 selected from the group consisting of 4-thiazolidinone, 5-[(2,3-dimethoxyphenyl)methylene]-2-thioxo-;
 4-thiazolidinone, 5-[(2,4-dimethoxyphenyl)methylene]-2-thioxo-;
 4-thiazolidinone, 2-thioxo-5-[(2,3,4-trimethoxyphenyl)methylene]-;
 4-thiazolidinone, 5-[(2,4-dichlorophenyl)methylene]-2-thioxo-;
 20 4-thiazolidinone, 5-[(3-fluoro-4-methoxyphenyl)methylene]-2-thioxo-;
 4-thiazolidinone, 5-[(3,4-dihydroxyphenyl)methylene]-2-thioxo-;
 4-thiazolidinone, 5-[(3-bromo-4-hydroxyphenyl)methylene]-2-thioxo-;
 4-thiazolidinone, 5-[(3,5-di-bromo-4-hydroxyphenyl)methylene]-2-thioxo-;
 5-[(phenyl)methylene]-2-thioxo-4-thiazolidinone;
 25 4-thiazolidinone, 5-[(2-methoxyphenyl)methylene]-2-thioxo-;
 4-thiazolidinone, 5-[(3-methoxyphenyl)methylene]-2-thioxo-;
 4-thiazolidinone, 5-[(3-hydroxyphenyl)methylene]-2-thioxo-;
 4-thiazolidinone, 5-[(4-hydroxyphenyl)methylene]-2-thioxo-;
 4-thiazolidinone, 5-[(3-ethoxyphenyl)methylene]-2-thioxo-;
 30 4-thiazolidinone, 5-[(3-(1-methylethoxy)phenyl)methylene]-2-thioxo-;
 4-thiazolidinone, 5-[(3-(phenylmethoxy)phenyl)methylene]-2-thioxo-;
 4-thiazolidinone, 5-[(3-(methylphenyl)methylene]-2-thioxo-;
 4-thiazolidinone, 2-thioxo-5-[(3-(trifluoromethyl)phenyl)methylene]-;
 4-thiazolidinone, 5-[(3-chlorophenyl)methylene]-2-thioxo-;
 35 4-thiazolidinone, 5-[(3-bromophenyl)methylene]-2-thioxo-;
 4-thiazolidinone, 5-[(3-nitrophenyl)methylene]-2-thioxo-;
 4-thiazolidinone, 5-[(4-chlorophenyl)methylene]-2-thioxo-;
 4-thiazolidinone, 5-[(4-bromophenyl)methylene]-2-thioxo-;
 4-thiazolidinone, 5-[(4-phenoxyphenyl)methylene]-2-thioxo-;
 40 4-thiazolidinone, 5-[(4-nitrophenyl)methylene]-2-thioxo-;
 4-thiazolidinone, 5-[(methylthiophenyl)methylene]-2-thioxo-;
 4-thiazolidinone, 5-[(4-hydroxy-3-methoxyphenyl)methylene]-2-thioxo- and disodium salt thereof;
 4-thiazolidinone, 5-[(4-methylphenyl)methylene]-2-thioxo-;
 4-thiazolidinone, 5-[(1,1'-biphenyl)-4-ylmethylene]-2-thioxo-;
 45 4-thiazolidinone, 5-[(2,6-dichlorophenyl)methylene]-2-thioxo-;
 4-thiazolidinone, 5-[(3-fluorophenyl)methylene]-2-thioxo-;
 4-thiazolidinone, 5-[(2-hydroxyphenyl)methylene]-2-thioxo-;
 4-thiazolidinone, 5-[(3-hydroxy-4,5-dimethoxyphenyl)methylene]-2-thioxo-;
 4-thiazolidinone, 5-[(2,6-dimethoxyphenyl)methylene]-2-thioxo-;
 50 4-thiazolidinone, 5-[(2-hydroxy-5-methoxyphenyl)methylene]-2-thioxo-;
 4-thiazolidinone, 5-[(3-hydroxy-5-methoxyphenyl)methylene]-2-thioxo-;
 4-thiazolidinone, 5-[(3-(1,1-dimethylethyl)-4-hydroxy-5-methoxyphenyl)methylene]-;
 4-thiazolidinone, 5-[(3-[(dimethylamino)methyl]-4-hydroxy-5-methoxyphenyl)methylene]-2-thioxo-; and
 4-thiazolidinone, 5-[(2-hydroxy-4,5-dimethoxyphenyl)methylene]-2-thioxo-.
12. A composition of Claim 1 selected from the group consisting of 4-thiazolidinone, 5-[(5-bromo-2-thienyl)methylene]-2-thioxo-;
 4-thiazolidinone, 5-[(5-bromo-2-thienyl)methylene]-2-thioxo-, monosodium salt;
 4-thiazolidinone, 5-[(6-methoxy-2-benzofuranyl)methylene]-2-thioxo-;

- 4-thiazolidinone, 5-[(5-methoxy-2-benzofuranyl)methylene]-2-thio-;
 5-[(benzo[b]thien-2-yl)methylene]-2-thio-4-thiazolidinone;
 4-thiazolidinone, 5-[[5-methoxy-3-(1-methylethoxy)benzo[b]thien-2-yl]methylene]-2-thio-;
 4-thiazolidinone, 5-[(5-methoxy-1-phenyl-1H-indol-2-yl)methylene]-2-thio-;
 5 4-thiazolidinone, 5-[(5-methoxy-1-phenyl-1H-indol-2-yl)methylene]-2-thio, 1-methylpiperidine salt;
 5-[(4-pyridyl)methylene]-2-thio-4-thiazolidinone;
 5-[(2-thienyl)methylene]-2-thio-4-thiazolidinone;
 5-[(5-nitro-2-furanyl)methylene]-2-thio-4-thiazolidinone; and
 5-[(1-phenyl-1H-indol-2-yl)methylene]-2-thio-4-thiazolidinone.
- 10 13. A composition of Claim 4 which is 5-[(4-hydroxy-3,5-dimethoxyphenyl)methylene]-2,4-thiazolidinedione or
 5-[(4-hydroxy-3-methoxyphenyl)methylene]-2,4-thiazolidinedione.
14. A composition of Claim 4 selected from the group consisting of 5-[(3,5-bis(1,1-dimethylethyl)-4-hydroxyphenyl)methylene]-2,4-thiazolidinedione and its choline salt;
 15 5-[(phenyl)methylene]-2,4-thiazolidinedione;
 2,4-thiazolidinedione, 5-(m-methoxybenzylidene-; and
 5-[(4-methoxyphenyl)methylene]-2,4-thiazolidinedione.
15. A composition of Claim 5 which is 5-[(4-hydroxy-3-methoxyphenyl)methylene]-2-thio-4-oxazolidinone or
 20 5-[(4-hydroxy-3,5-dimethoxyphenyl)methylene]-2-thio-4-oxazolidinone and its disodium salt.
16. A composition of Claim 5 selected from the group consisting of 4-oxazolidinone, 5-[(3-methoxyphenyl)methylene]-2-thio- and its disodium salt;
 4-oxazolidinone, 5-[(4-methoxyphenyl)methylene]-2-thio-;
 4-oxazolidinone, 5-[(3,4-dimethoxyphenyl)methylene]-2-thio-;
 25 4-oxazolidinone, 5-[(4-hydroxy-3,5-dimethylphenyl)methylene]-2-thio-;
 4-oxazolidinone, 5-[(3-bromo-4-hydroxy-5-methoxyphenyl)methyl]-2-thio-;
 4-oxazolidinone, 2-thio-5-[(2,4,5-trimethoxyphenyl)methylene]-; and
 4-oxazolidinone, 2-thio-5-[(3-bromo-4-hydroxyphenyl)methylene]-.
- 30 17. A composition of Claim 5 which is 2-thio-5-[(3,4,5-trimethoxyphenyl)methylene]-4-oxazolidinone.
18. A composition of Claim 6 which is 5-[(4-hydroxy-3,5-dimethoxyphenyl)methylene]-2-thio-4-imidazolidinone.
19. A composition of Claim 6 selected from the group consisting of 4-imidazolidinone, 5-[(3,5-dibromo-4-hydroxyphenyl)methylene]-2-thio-;
 35 4-imidazolidinone, 5-[(3-bromo-4-hydroxyphenyl)methylene]-2-thio-;
 4-imidazolidinone, 5-[[3-bis(1,1-dimethylethyl)-4-hydroxyphenyl]methylene]-2-thio-;
 4-imidazolidinone, 5-[2-methoxyphenyl)methylene]-2-thio-;
 4-imidazolidinone, 5-[(3-methoxyphenyl)methylene]-2-thio-;
 4-imidazolidinone, 5-[(4-methoxyphenyl)methylene]-2-thio-;
 40 Hydantoin, 2-thio-5-vanillylidene-; and
 4-imidazolidinone, 5-[(4-hydroxy-3,5-dimethoxyphenyl)methylene]-2-thio-.
20. A composition of Claim 7 selected from the group consisting of 2,4-imidazolidinedione, 5-[[4-(acetyloxy)-3-methoxyphenyl)methylene]-;
 2,4-imidazolidinedione, 5-[(4-hydroxy-3-methoxyphenyl)methylene]-;
 45 2,4-imidazolidinedione, 5-[(4-hydroxy-3,5-dimethoxyphenyl)methylene]-;
 2,4-imidazolidinedione, 5-[[3,5-bis(1,1-dimethylethyl)-4-hydroxyphenyl)methylene]-;
 2,4-imidazolidinedione, 5-[(3-methoxyphenyl)methylene]-1-methyl-;
 2,4-imidazolidinedione, 5-[(4-hydroxy-3-methoxyphenyl)methylene]-1-methyl-;
 2,4-imidazolidinedione, 5-[(4-hydroxy-3,5-dimethoxyphenyl)methylene]-1-methyl-;
 50 2,4-imidazolidinedione, 5-[(3,5-dibromo-4-hydroxyphenyl)methylene]-1-methyl-; and
 2,4-imidazolidinedione, 5-[(4-hydroxy-3,5-bis(1-methylethyl)phenyl)methylene]-1-methyl-.
21. A composition of Claim 8 selected from the group consisting of 2,4-imidazolidinedione, 5-[(4-methoxyphenyl)methylene]-3-methyl-;
 2,4-imidazolidinedione, 5-[(3-methoxyphenyl)methylene]-3-methyl-;
 55 2,4-imidazolidinedione, 5-[(4-methoxyphenyl)methylene]-3-methyl-;
 Hydantoin, 3-methyl-5-vanillylidene-; and
 2,4-imidazolidinedione, 5-[(4-hydroxy-3,5-dimethoxyphenyl)methylene]-3-methyl-.

22. A composition of Claim 9 selected from the groups consisting of
 5-[[3,5-bis(1,1-dimethylethyl)-4-hydroxyphenyl]methylene]-2,4-oxazolidinone;
 5-[(4-hydroxy-3,5-dimethoxyphenyl)methylene]-2,4-oxazolidinone; and
 5-[(4-hydroxy-3,5-dimethylphenyl)methylene]-2,4-oxazolidinone.

23. A compound selected from the group consisting of
 5-[(3-fluoro-4-methoxyphenyl)methylene]-2-thioxo-4-thiazolidinone;
 5-[(3-bromo-4-hydroxyphenyl)methylene]-2-thioxo-4-thiazolidinone;
 5-[(3,5-dibromo-4-hydroxyphenyl)methylene]-2-thioxo-4-thiazolidinone; and its disodium salt;
 5-[(3-bromo-4-hydroxy-5-methoxyphenyl)methylene]-2-thioxo-4-thiazolidinone and its disodium salt;
 5-[[3,5-bis(1,1-dimethylethyl)-4-hydroxyphenyl]methylene]-2-thioxo-oxazolidinone and its choline salt.
 5-[(3-fluorophenyl)methylene]-2-thioxo-4-thiazolidinone;
 5-[(4-hydroxy-3-methoxy-5-tertiarybutylphenyl)methylene]-2-thioxo-4-thiazolidinone;
 5-[2-hydroxy-4,5-dimethoxyphenyl)methylene]-2-thioxo-4-thiazolidinone;
 5-[[3-[(dimethylamino)methylene]-2-thioxo-4-thiazolidinone;
 5-[(4-hydroxy-3,5-dimethoxyphenyl)methylene]-2,4-thiazolidinedione;
 5-[(3-methoxyphenyl)methylene]-2-thioxo-4-oxazolidinone;
 5-[(3,4-dimethoxyphenyl)methylene]-2-thioxo-4-oxazolidinone;
 5-[(4-hydroxy-3,5-dimethoxyphenyl)methylene]-2-thioxo-4-oxazolidinone;
 5-[(4-hydroxy-3,5-dimethylphenyl)methylene]-2-thioxo-4-oxazolidinone;
 5-[(3-bromo-4-hydroxy-5-methoxyphenyl)methylene]-2-thioxo-4-oxazolidinone;
 2-thioxo-5-[(2,4,5-trimethoxyphenyl)methylene]-4-oxazolidinone;
 5-[(3,5-dibromo-4-hydroxyphenyl)methylene]-2-thioxo-4-imidazolidinone;
 5-[(3-bromo-4-hydroxyphenyl)methylene]-2-thioxo-4-imidazolidinone;
 5-[(3,5-dimethoxy-4-hydroxyphenyl)methylene]-2-thioxo-4-imidazolidinone;
 5-[(3,5-dimethoxy-4-hydroxyphenyl)methylene]-3-methyl-2,4-imidazolidinedione;
 5-[(4-hydroxy-3-methoxyphenyl)methylene]-1-methyl-2,4-imidazolidinedione;
 5-[(3,5-dimethoxy-4-hydroxyphenyl)methylene]-1-methyl-2,4-imidazolidinedione;
 5-[(3,5-dibromo-4-hydroxyphenyl)methylene]-1-methyl-2,4-imidazolidinedione;
 5-[[3,5-bis(1,1-dimethylethyl)-4-hydroxyphenyl]methylene]-2,4-thiazolidinedione;
 5-[[3,5-bis(1,1-dimethylethyl)-4-hydroxyphenyl]methylene]-2-thioxo-4-oxazolidinone;
 5-[[3,5-bis(1,1-dimethylethyl)-4-hydroxyphenyl]methylene]-2,4-oxazolidinone;
 5-[(4-hydroxy-3,5-dimethoxyphenyl)methylene]-2,4-oxazolidinedione;
 5-[(4-hydroxy-3,5-dimethylphenyl)methylene]-2,4-oxazolidinone.
 24. A compound selected from the group consisting of 4-thiazolidinone, 5-[(6-methoxy-2-benzofuranyl)methylene]-2-thioxo-;
 4-thiazolidinone, 5-[(5-methoxy-2-benzofuranyl)methylene]-2-thioxo-;
 Rhodanine, 5-(benzo[b]thien-2-ylmethylene)-;
 4-thiazolidinone, 5-[[5-methoxy-3-(1-methylethoxy)benzo[b]thien-2-yl]methylene]-2-thioxo-; and
 4-thiazolidinone, 5-[[5-methoxy-1-phenyl-1H-indol-2-yl]methylene]-2-thioxo-.
25. A use of a compound of the formula (I)



- and pharmaceutically acceptable salts thereof;
 wherein Ar is (i) phenyl;
 (ii) phenyl substituted by from one to three of lower alkyl, lower alkoxy, hydroxy, halogen, trifluoromethyl, NO₂, mercapto, or lower alkylthio, NR₄R₅ wherein R₄ and R₅ are independently hydrogen, or lower alkyl;
 (iii) naphthyl;
 (iv) benzofuranyl;
 (v) benzothiophenyl;
 (vi) thienyl;
 (vii) indolyl

(viii) furanyl

(ix) pyridyl

Y and Y₁ is oxygen or sulfur;

X is sulfur, oxygen, NH or NCH₃;


5 X₁ is NH or NCH₃;

with the proviso that Ar is not 3,5-di(isopropyl)-4-hydroxyphenyl and further if X₁ is NH, X is S, and Y is S then Ar is not 3,5-di(t-butyl)-4-hydroxyphenyl and a pharmaceutically acceptable carrier in the manufacture of medical therapeutic agents for treating allergies and inflammations.

26. A pharmaceutical composition as claimed in Claim 1 additionally comprising an effective amount of
 10 a second active ingredient that is a nonsteroidal antiinflammatory drug; a peripheral analgesic agent; a cyclooxygenase inhibitor; a leukotriene antagonist; an antihistaminic agent; a prostaglandin antagonist or a thromboxane antagonist.

15 Claims for the following Contracting States: ES, GR

1. A process for preparing a pharmaceutical composition comprising the addition of a compound of the formula (I)



and pharmaceutically acceptable salts thereof;

wherein Ar is (i) phenyl;

(ii) phenyl substituted by from one to three of lower alkyl, lower alkoxy, hydroxy, halogen, trifluoromethyl,
 30 NO₂, mercapto, or lower alkylthio, NR₁₀R₁₁ wherein R₁₀ and R₁₁ are independently hydrogen, or lower alkyl;

(iii) naphthyl;

(iv) benzofuranyl;

(v) benzothiophenyl;

35 (vi) thienyl;

(vii) indolyl

(viii) furanyl

(ix) pyridyl


Y and Y₁ is oxygen or sulfur;

40 X is sulfur, oxygen, NH or NCH₃; and

X₁ is NH or NCH₃;


with the proviso that Ar is not 3,5-di(isopropyl)-4-hydroxyphenyl and further if X₁ is NH, X is S, and Y is S then Ar is not 3,5-di(t-butyl)-4-hydroxyphenyl to a pharmaceutically acceptable carrier.

2. A process of Claim 1 wherein the compound is of the formula (II)



wherein X and Y are as defined above and having of from one to three of R₁, R₂, R₃, R₄ and R₅ selected
 55 from the group consisting of lower alkyl, lower alkoxy, OH, halogen, CF₃, NR₁₀R₁₁ wherein R₁₀ and R₁₁ are as defined above, NO₂, mercapto, or lower alkylthio.


3. A process of Claim 2 wherein the compound is of the formula (III)



III


wherein R_1 , R_2 , R_3 , R_4 and R_5 are as defined above.

4. A process of Claim 2 wherein the compound is of the formula (IV)



IV


5. A process of Claim 2 wherein the compound is of the formula (V)



V

wherein R_1 , R_2 , R_3 , R_4 and R_5 are as defined above.


6. A process of Claim 2 wherein the compound is of the formula (VI)



VI

wherein R_1 , R_2 , R_3 , R_4 and R_5 are as defined above.


7. A process of Claim 2 wherein the compound is of the formula (VII)



VII

wherein R_1 , R_2 , R_3 , R_4 and R_5 are as defined above.


8. A process of Claim 1 wherein the compound is of the formula (VIII)



VIII

wherein R₁, R₂, R₃, R₄ and R₅ are as defined above.

9. A process of Claim 2 wherein the compound is of the formula (VIII A)



VIII A

wherein R₁, R₂, R₃, R₄, and R₅ are as defined above.

10. A process of Claim 3 wherein the compound is a member of the group consisting of

5-[(2,5-dimethoxyphenyl)methylene]-2-thioxo-4-thiazolidinone;
 5-[(2,4,5-trimethoxyphenyl)methylene]-2-thioxo-4-thiazolidinone;
 the monosodium salt of 5-[(2,4,5-trimethoxyphenyl)methylene]-2-thioxo-4-thiazolidinone;
 5-[(3,4,5-trimethoxyphenyl)methylene]-2-thioxo-4-thiazolidinone;
 5-[(3,4-dichlorophenyl)methylene]-2-thioxo-4-thiazolidinone;
 the monosodium salt of 5-[(3,4,5-trimethoxyphenyl)methylene]-2-thioxo-4-thiazolidinone;
 5-[(3,4-dimethoxyphenyl)methylene]-2-thioxo-4-thiazolidinone;
 the monosodium salt of 5-[(3,4-dichlorophenyl)methylene]-2-thioxo-4-thiazolidinone;
 5-[(3,5-dimethoxy-4-hydroxyphenyl)methylene]-2-thioxo-4-thiazolidinone;
 the disodium salt of 5-[(3,4,5-trimethoxyphenyl)methylene]-2-thioxo-4-thiazolidinone;
 5-[(3,5-dimethyl-4-hydroxyphenyl)methylene]-2-thioxo-4-thiazolidinone;
 the disodium salt of 5-[(3,4-dimethoxyphenyl)methylene]-2-thioxo-4-thiazolidinone;
 5-[(3-bromo-4-hydroxy-5-methoxyphenyl)methylene]-2-thioxo-4-thiazolidinone;
 the disodium salt of 5-[(3,4-dichlorophenyl)methylene]-2-thioxo-4-thiazolidinone;
 5-[(4-methoxyphenyl)methylene]-2-thioxo-4-thiazolidinone;
 5-[(3-hydroxy-4-methoxyphenyl)methylene]-2-thioxo-4-thiazolidinone;
 5-[(3,5-dimethoxyphenyl)methylene]-2-thioxo-4-thiazolidinone.

11. A process of Claim 3 wherein the compound is selected from the group consisting of 4-

thiazolidinone, 5-[(2,3-dimethoxyphenyl)methylene]-2-thioxo-;
 4-thiazolidinone, 5-[(2,4-dimethoxyphenyl)methylene]-2-thioxo-;
 4-thiazolidinone, 2-thioxo-5-[(2,3,4-trimethoxyphenyl)methylene]-;
 4-thiazolidinone, 5-[(2,4-dichlorophenyl)methylene]-2-thioxo-;
 4-thiazolidinone, 5-[(3-fluoro-4-methoxyphenyl)methylene]-2-thioxo-;
 4-thiazolidinone, 5-[(3,4-dihydroxyphenyl)methylene]-2-thioxo-;
 4-thiazolidinone, 5-[(3-bromo-4-hydroxyphenyl)methylene]-2-thioxo-;
 4-thiazolidinone, 5-[(3,5-di-bromo-4-hydroxyphenyl)methylene]-2-thioxo-;
 5-[(phenyl)methylene]-2-thioxo-4-thiazolidinone;
 4-thiazolidinone, 5-[(2-methoxyphenyl)methylene]-2-thioxo-;
 4-thiazolidinone, 5-[(3-methoxyphenyl)methylene]-2-thioxo-;
 4-thiazolidinone, 5-[(3-hydroxyphenyl)methylene]-2-thioxo-;
 4-thiazolidinone, 5-[(4-hydroxyphenyl)methylene]-2-thioxo-;
 4-thiazolidinone, 5-[(3-ethoxyphenyl)methylene]-2-thioxo-;
 4-thiazolidinone, 5-[(3-(1-methylethoxy)phenyl)methylene]-2-thioxo-;
 4-thiazolidinone, 5-[(3-(phenylmethoxy)phenyl)methylene]-2-thioxo-;
 4-thiazolidinone, 5-[(3-methylphenyl)methylene]-2-thioxo-;
 4-thiazolidinone, 2-thioxo-5-[(3-trifluoromethyl)phenyl)methylene]-;
 4-thiazolidinone, 5-[(3-chlorophenyl)methylene]-2-thioxo-;

- 4-thiazolidinone, 5-[(3-bromophenyl)methylene]-2-thioxo-;
 4-thiazolidinone, 5-[(3-nitrophenyl)methylene]-2-thioxo-;
 4-thiazolidinone, 5-[(4-chlorophenyl)methylene]-2-thioxo-;
 4-thiazolidinone, 5-[(4-bromophenyl)methylene]-2-thioxo-;
 5 4-thiazolidinone, 5-[(4-phenoxyphenyl)methylene]-2-thioxo-;
 4-thiazolidinone, 5-[(4-nitrophenyl)methylene]-2-thioxo-;
 4-thiazolidinone, 5-[(4-methylthiophenyl)methylene]-2-thioxo-;
 4-thiazolidinone, 5-[(4-hydroxy-3-methoxyphenyl)methylene]-2-thioxo- and disodium salt thereof;
 4-thiazolidinone, 5-[(4-methylphenyl)methylene]-2-thioxo-;
 10 4-thiazolidinone, 5-[(1,1'-biphenyl)-4-ylmethylene]-2-thioxo-;
 4-thiazolidinone, 5-[(2,6-dichlorophenyl)methylene]-2-thioxo-;
 4-thiazolidinone, 5-[(3-fluorophenyl)methylene]-2-thioxo-;
 4-thiazolidinone, 5-[(2-hydroxyphenyl)methylene]-2-thioxo-;
 4-thiazolidinone, 5-[(3-hydroxy-4,5-dimethoxyphenyl)methylene]-2-thioxo-;
 15 4-thiazolidinone, 5-[(2,6-dimethoxyphenyl)methylene]-2-thioxo-;
 4-thiazolidinone, 5-[(2-hydroxy-5-methoxyphenyl)methylene]-2-thioxo-;
 4-thiazolidinone, 5-[(3-hydroxy-5-methoxyphenyl)methylene]-2-thioxo-;
 4-thiazolidinone, 5-[(3-(1,1-dimethylethyl)-4-hydroxy-5-methoxyphenyl)methylene]-;
 4-thiazolidinone, 5-[(3-[(dimethylamino)methyl]-4-hydroxy-5-methoxyphenyl)methylene]-2-thioxo-; and
 20 4-thiazolidinone, 5-[(2-hydroxy-4,5-dimethoxyphenyl)methylene]-2-thioxo-.
12. A process of Claim 1 wherein the compound is selected from the group consisting of 4-thiazolidinone, 5-[(5-bromo-2-thienyl)methylene]-2-thioxo-;
 4-thiazolidinone, 5-[(5-bromo-2-thienyl)methylene]-2-thioxo-, monosodium salt;
 4-thiazolidinone, 5-[(6-methoxy-2-benzofuranyl)methylene]-2-thioxo-;
 25 4-thiazolidinone, 5-[(5-methoxy-2-benzofuranyl)methylene]-2-thioxo-;
 5-[(benzo[b]thien-2-yl)methylene]-2-thioxo-4-thiazolidinone;
 4-thiazolidinone, 5-[(5-methoxy-3-(1-methylethoxy)benzo[b]thien-2-yl)methylene]-2-thioxo-;
 4-thiazolidinone, 5-[(5-methoxy-1-phenyl-1H-indol-2-yl)methylene]-2-thioxo-;
 4-thiazolidinone, 5-[(5-methoxy-1-phenyl-1H-indol-2-yl)methylene]-2-thioxo, 1-methylpiperidine salt;
 30 5-[(4-pyridyl)methylene]-2-thioxo-4-thiazolidinone;
 5-[(2-thienyl)methylene]-2-thioxo-4-thiazolidinone;
 5-[(5-nitro-2-furanyl)methylene]-2-thioxo-4-thiazolidinone; and
 5-[(1-phenyl-1H-indol-2-yl)methylene]-2-thioxo-4-thiazolidinone.
13. A process of Claim 4 wherein the compound is 5-[(4-hydroxy-3,5-dimethoxyphenyl)methylene]-2,4-thiazolidinedione or
 35 5-[(4-hydroxy-3-methoxyphenyl)methylene]-2,4-thiazolidinedione.
14. A process of Claim 4 wherein the compound is selected from the group consisting of 5-[(3,5-bis(1,1-dimethylethyl)-4-hydroxyphenyl)methylene]-2,4-thiazolidinedione and its choline salt;
 5-[(phenyl)methylene]-2,4-thiazolidinedione;
 40 2,4-thiazolidinedione, 5-(m-methoxybenzylidene)-; and
 5-[(4-methoxyphenyl)methylene]-2,4-thiazolidinedione.
15. A process of Claim 5 wherein the compound is 5-[(4-hydroxy-3-methoxyphenyl)methylene]-2-thioxo-4-oxazolidinone or
 5-[(4-hydroxy-3,5-dimethoxyphenyl)methylene]-2-thioxo-4-oxazolidinone and its disodium salt.
 45 16. A process of Claim 5 wherein the compound is selected from the group consisting of 4-oxazolidinone, 5-[(3-methoxyphenyl)methylene]-2-thioxo- and its disodium salt;
 4-oxazolidinone, 5-[(4-methoxyphenyl)methylene]-2-thioxo-;
 4-oxazolidinone, 5-[(3,4-dimethoxyphenyl)methylene]-2-thioxo-;
 4-oxazolidinone, 5-[(4-hydroxy-3,5-dimethylphenyl)methylene]-2-thioxo-;
 50 4-oxazolidinone, 5-[(3-bromo-4-hydroxy-5-methoxyphenyl)methyl]-2-thioxo-;
 4-oxazolidinone, 2-thioxo-5-[(2,4,5-trimethoxyphenyl)methylene]-; and
 4-oxazolidinone, 2-thioxo-5-[(3-bromo-4-hydroxyphenyl)methylene]-.
17. A process of Claim 5 wherein the compound is 2-thioxo-5-[(3,4,5-trimethoxyphenyl)methylene]-4-oxazolidinone.
- 55 18. A process of Claim 6 wherein the compound is 5-[(4-hydroxy-3,5-dimethoxyphenyl)methylene]-2-thioxo-4-imidazolidinone.

19. A process of Claim 6 wherein the compound is selected from the group consisting of 4-imidazolidinone, 5-[(3,5-dibromo-4-hydroxyphenyl)methylene]-2-thioxo-; 4-imidazolidinone, 5-[(3-bromo-4-hydroxyphenyl)methylene]-2-thioxo-; 4-imidazolidinone, 5-[(3-bis(1,1-dimethylethyl)-4-hydroxyphenyl)methylene]-2-thioxo-; 4-imidazolidinone, 5-[2-methoxyphenyl)methylene]-2-thioxo-; 4-imidazolidinone, 5-[(3-methoxyphenyl)methylene]-2-thioxo-; 4-imidazolidinone, 5-[(4-methoxyphenyl)methylene]-2-thioxo-; Hydantoin, 2-thio-5-vanillylidene-; and 4-imidazolidinone, 5-[(4-hydroxy-3,5-dimethoxyphenyl)methylene]-2-thioxo-.
20. A process of Claim 7 wherein the compound is selected from the group consisting of 2,4-imidazolidinedione, 5-[(4-acetyloxy-3-methoxyphenyl)methylene]-; 2,4-imidazolidinedione, 5-[(4-hydroxy-3-methoxyphenyl)methylene]-; 2,4-imidazolidinedione, 5-[(4-hydroxy-3,5-dimethoxyphenyl)methylene]-; 2,4-imidazolidinedione, 5-[(3,5-bis(1,1-dimethylethyl)-4-hydroxyphenyl)methylene]-; 2,4-imidazolidinedione, 5-[(3-methoxyphenyl)methylene]-1-methyl-; 2,4-imidazolidinedione, 5-[(4-hydroxy-3-methoxyphenyl)methylene]-1-methyl-; 2,4-imidazolidinedione, 5-[(4-hydroxy-3,5-dimethoxyphenyl)methylene]-1-methyl-; 2,4-imidazolidinedione, 5-[(3,5-dibromo-4-hydroxyphenyl)methylene]-1-methyl-; and 2,4-imidazolidinedione, 5-[(4-hydroxy-3,5-bis(1-methylethyl)phenyl)methylene]-1-methyl-.
21. A process of Claim 8 wherein the compound is selected from the group consisting of 2,4-imidazolidinedione, 5-[(2-methoxyphenyl)methylene]-3-methyl-; 2,4-imidazolidinedione, 5-[(3-methoxyphenyl)methylene]-3-methyl-; 2,4-imidazolidinedione, 5-[(4-methoxyphenyl)methylene]-3-methyl-; Hydantoin, 3-methyl-5-vanillylidene-; and 2,4-imidazolidinedione, 5-[(4-hydroxy-3,5-dimethoxyphenyl)methylene]-3-methyl-.
22. A process of Claim 9 wherein the compound is selected from the group consisting of 5-[(3,5-bis(1,1-dimethylethyl)-4-hydroxyphenyl)methylene]-2,4-oxazolidinone; 5-[(4-hydroxy-3,5-dimethoxyphenyl)methylene]-2,4-oxazolidinone; and 5-[(4-hydroxy-3,5-dimethylphenyl)methylene]-2,4-oxazolidinone.
23. A process for the preparation of a compound according to Claim 1, namely a member selected from the group consisting of 5-[(3-fluoro-4-methoxyphenyl)methylene]-2-thioxo-4-thiazolidinone; 5-[(3-bromo-4-hydroxyphenyl)methylene]-2-thioxo-4-thiazolidinone; 5-[(3,5-dibromo-4-hydroxyphenyl)methylene]-2-thioxo-4-thiazolidinone; and its disodium salt; 5-[(3-bromo-4-hydroxy-5-methoxyphenyl)methylene]-2-thioxo-4-thiazolidinone and its disodium salt; 5-[(3,5-bis(1,1-dimethylethyl)-4-hydroxyphenyl)methylene]-2-thioxo-oxazolidinone and its choline salt; 5-[(3-fluorophenyl)methylene]-2-thioxo-4-thiazolidinone; 5-[(4-hydroxy-3-methoxy-5-tertiarybutylphenyl)methylene]-2-thioxo-4-thiazolidinone; 5-[2-hydroxy-4,5-dimethoxyphenyl)methylene]-2-thioxo-4-thiazolidinone; 5-[(3-[(dimethylamino)methylene]-2-thioxo-4-thiazolidinone; 5-[(4-hydroxy-3,5-dimethoxyphenyl)methylene]-2,4-thiazolidinedione; 5-[(3-methoxyphenyl)methylene]-2-thioxo-4-oxazolidinone; 5-[(3,4-dimethoxyphenyl)methylene]-2-thioxo-4-oxazolidinone; 5-[(4-hydroxy-3,5-dimethoxyphenyl)methylene]-2-thioxo-4-oxazolidinone; 5-[(4-hydroxy-3,5-dimethylphenyl)methylene]-2-thioxo-4-oxazolidinone; 5-[(3-bromo-4-hydroxy-5-methoxyphenyl)methylene]-2-thioxo-4-oxazolidinone; 2-thioxo-5-[(2,4,5-trimethoxyphenyl)methylene]-4-oxazolidinone; 5-[(3,5-dibromo-4-hydroxyphenyl)methylene]-2-thioxo-4-imidazolidinone; 5-[(3-bromo-4-hydroxyphenyl)methylene]-2-thioxo-4-imidazolidinone; 5-[(3,5-dimethoxy-4-hydroxy)methylene]-2-thioxo-4-imidazolidinone; 5-[(3,5-dimethoxy-4-hydroxyphenyl)methylene]-3-methyl-2,4-imidazolidinedione; 5-[(4-hydroxy-3-methoxyphenyl)methylene]-1-methyl-2,4-imidazolidinedione; 5-[(3,5-dimethoxy-4-hydroxyphenyl)methylene]-1-methyl-2,4-imidazolidinedione; 5-[(3,5-dibromo-4-hydroxyphenyl)methylene]-1-methyl-2,4-imidazolidinedione; 5-[(3,5-bis(1,1-dimethylethyl)-4-hydroxyphenyl)methylene]-, 2,4-thiazolidinedione; 5-[(3,5-bis(1,1-dimethylethyl)-4-hydroxyphenyl)methylene]-2-thioxo-4-oxazolidinone; 5-[(3,5-bis(1,1-dimethylethyl)-4-hydroxyphenyl)methylene]-2,4-oxazolidinone; 5-[(4-hydroxy-3,5-dimethoxyphenyl)methylene]-2,4-oxazolidinedione;

5-[(4-hydroxy-3,5-dimethylphenyl)methylene]-2,4-oxazolidinone;

4-thiazolidinone, 5-[(6-methoxy-2-benzofuranyl)methylene]-2-thio-


4-thiazolidinone, 5-[(5-methoxy-2-benzofuranyl)methylene]-2-thio-

Rhodanine, 5-(benzo[b]thien-2-yl)methylene-

5 4-thiazolidinone, 5-[[5-methoxy-3-(1-methylethoxy)benzo[b]thien-2-yl]methylene]-2-thio-; and


4-thiazolidinone, 5-[(5-methoxy-1-phenyl-1H-indol-2-yl)methylene]-2-thio-

characterized by the aldol-condensation of an aldehyde of the formula ArCHO , wherein Ar has the meaning of Claim 1, with a compound of formula



wherein X, X₁, Y, Y₁ have the meaning of Claim 1, whereafter the compound obtained is optionally hydrolysed and/or transferred to a pharmaceutically acceptable salt.

24. A use of a compound of the formula (I)



and pharmaceutically acceptable salts thereof;

wherein Ar is (i) phenyl;

(ii) phenyl substituted by from one to three of lower alkyl, lower alkoxy, hydroxy, halogen, trifluoromethyl, NO₂, mercapto, or lower alkylthio, NR₄R₅ wherein R₄ and R₅ are independently hydrogen, or lower alkyl;

(iii) naphthyl;

(iv) benzofuranyl;

(v) benzothiophenyl;

(vi) thienyl;

(vii) indolyl

(viii) furanyl

(ix) pyridyl

Y and Y₁ is oxygen or sulfur;

X is sulfur, oxygen, NH or NCH₃;

X₁ is NH or NCH₃;

with the proviso that Ar is not 3,5-di(isopropyl)-4-hydroxyphenyl and further if X₁ is NH, X is S, and Y is S then Ar is not 3,5-di(t-butyl)-4-hydroxyphenyl and a pharmaceutically acceptable carrier in the manufacture

of medical therapeutic agents for treating allergies and inflammations.

THIS PAGE BLANK (USPTO)

12

EUROPEAN PATENT APPLICATION

21 Application number: 89109406.2

22 Date of filing: 24.05.89

51 Int. Cl. 4: A 61 K 31/41

A 61 K 31/415, A 61 K 31/42,
 A 61 K 31/425, A 61 K 31/44,
 A 61 K 31/33, A 61 K 45/06,
 C 07 D 233/66,
 C 07 D 277/32,
 C 07 D 263/34, C 07 D 417/06

30 Priority: 25.05.88 US 198528 10.04.89 US 334346

43 Date of publication of application:
29.11.89 Bulletin 89/48

84 Designated Contracting States:
AT BE CH DE ES FR GB GR IT LI LU NL SE

88 Date of deferred publication of search report:
21.03.90 Bulletin 90/12

71 Applicant: WARNER-LAMBERT COMPANY
201 Tabor Road
Morris Plains New Jersey 07950 (US)

72 Inventor: Cetenko, Wlaczslaw Antin
3565 Green Brier Road
Ann Arbor Michigan 48105 (US)

Connor, David Thomas
 2453 Antietam
 Ann Arbor Michigan 48105 (US)

Sorenson, Roderick Joseph
 2820 Briarcliff
 Ann Arbor Michigan 48105 (US)


Unangst, Paul Charles
 3659 Middleton Drive
 Ann Arbor Michigan 48105 (US)

Stabler, Stephen Russell
 2211 Latham Street No. 304
 Mountainview California 94040 (US)

74 Representative: Mansmann, Ivo
 c/o Gödecke AG - Patentabteilung Postfach 569
 Mooswaldallee 1-9
 D-7800 Freiburg (DE)

54 Known and selected novel arylmethylenyl derivatives of thiazolidinones, imidazolidinones and oxazolidinones useful as antiallergy agents and antiinflammatory agents.

57 The present invention is for selected novel compounds, as well as, pharmaceutical compositions and methods of use for known and the selected novel compounds both of the formula



having activity useful for treating allergies and inflammation.

EP 0 343 643 A3



DOCUMENTS CONSIDERED TO BE RELEVANT			EP 89109406.2
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl. 4)
D,X	J. PHARM. SOC. JAP., vol. 76, 1956 H.TANIYAMA et al. "Studies on Chemotherapeutics for Mycobacterium tuberculosis. XI. Synthesis and Antibacterial Activity of 4-Thiazolidine-Derivatives containing the alpha,beta-Unsaturated Ketone Group (1)" pages 154-157 * Abstract; table I *	1-4, 11,14	A 61 K 31/41 A 61 K 31/415 A 61 K 31/42 A 61 K 31/425 A 61 K 31/44 A 61 K 31/33 A 61 K 45/06 C 07 D 233/66 C 07 D 277/32 C 07 D 263/34 C 07 D 417/06
D,X	EGYPTIAN JOURNAL OF CHEMISTRY, vol. 26, 1983, The Egyptian Chemical Society H.H.MOHARRAM et al. "Synthesis of some New Rhodanine Derivatives of Antibacterial and Anticancer Effects" pages 301-311 * Page 301 (Ia,c,d), tables 1,3 *	1	TECHNICAL FIELDS SEARCHED (Int. Cl. 4)
D,X	CHEMICAL & PHARMACEUTICAL BULLETIN, vol. 34, 1986, Pharmaceutical Society of Japan J.KATSUMI et al. "Studies on Styrene Derivatives. II. Synthesis and Antiinflammatory Activity of 3,5-Di-tert-butyl-4-hydroxystyrenes" pages 1619-1627 * Table 1, compounds no. 11-12, page 1624, first passage after table 1 *	1-3,6, 7,25	A 61 K 31/00 C 07 D 233/00 C 07 D 277/00 C 07 D 263/00 C 07 D 417/00 C 07 D 209/00
D,X	INTERNATIONAL JOURNAL OF SULFUR CHEMISTRY, A-vol. 2,	23	
The present search report has been drawn up for all claims			
Place of search		Date of completion of the search	Examiner
VIENNA		15-11-1989	MAZZUCCO
CATEGORY OF CITED DOCUMENTS			
X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document		T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document	



DOCUMENTS CONSIDERED TO BE RELEVANT			CLASSIFICATION OF THE APPLICATION (Int. Cl. 4)
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	
X	no. 1, 1972, The Intra- -Science Research Foundation T.R.BOSIN et al. "A Novel Product From Oxidation of beta-(3-Malogenophenyl)- -alpha-Mercaptoacrylic Acids" pages 261-266 * Page 261, right column, second passage, compounds (1); page 262, formula 1; page 263, table 1, first formula * -- FARMATSEVTICHNIJ ZHURUAL, vol. 25, no. 6, 1970, Kiev N.E.PLEVACHUK "Synthesis and transformation of 3-alkyl (aryl)-4-thionothiazoli- dones" pages 3-6 * Page 4, first line right formula and table * (see also Chemical Abstracts 75, 5769r (1971)) --	1	
X	YAKUGAKU ZASSHI, vol. 92, 1972, Pharmaceutical Society of Japan M.GOSHA et al. "Antimicrobial Activity of 5-Oxothiazoli- dine-2-thione and Rhodanine Derivatives" pages 490-497 * Page 490, introduction, tables III, IV, V * (see also Chemical Abstracts 77, 70 744j (1972)) --	1-3	TECHNICAL FIELDS SEARCHED (Int. Cl. 4, --
D	EP - A2 - 0 211 670 (ELI LILLY AND COMPANY)		
The present search report has been drawn up for all claims			
Place of search		Date of completion of the search	Examiner
VIENNA		15-11-1989	MAZZUCCO
CATEGORY OF CITED DOCUMENTS			
X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document		T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document	



DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl. 4)
X	* Claims 1,3,4; page 3, line 30 - page 4, line 12; example 5; page 15, lines 23-30 *	1,2,25	
A	* Claims 1,3,4; page 3, line 30 - page 4, line 12; example 5; page 15, lines 23-30 *	3,4,11,23	
X	DE - A1 - 3 023 349 (DEGUSSA AG) * Page 6, line 6 - page 7, line 27 *	23	
D,A	PATENT ABSTRACTS OF JAPAN, unexamined applications, section C, vol. 10, no. 156, June 5, 1986 THE PATENT OFFICE JAPANESE GOVERNMENT page 126 C 351 * Kokai-no. 61-12 674 (KANEGAFUCHI KAGAKU KOGYO K.K.) *	1,2,7, 8,20	
A	ARZNEIMITTEL-FORSCHUNG, vol. 19, 1969, Editir Cantor KG T.ZSOLNAI "Die antimikro- bielle Wirkung von potentiellen Isothiocyanat- Bildnern" pages 558-572 * Page 561, compounds no. 110-115; page 570, point 14 *	1,2, 10,11, 23,25	
A	ACTA PHYTOPATHOLOGICA ACADEMIAE SCIENTARUM HUNGARICAE, vol. 4, 1969, Akademiai Kiado, Budapest	1,25	
The present search report has been drawn up for all claims			
Place of search VIENNA		Date of completion of the search 15-11-1989	Examiner MAZZUCCO
<p>CATEGORY OF CITED DOCUMENTS</p> <p>X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document</p> <p>T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document</p>			



European Patent
Office

EUROPEAN SEARCH REPORT

Application number

-4-

EP 89109406.2

DOCUMENTS CONSIDERED TO BE RELEVANT			CLASSIFICATION OF THE APPLICATION (Int. Cl.4)
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	
D	G.MATOLCSY et al. "Studies on the Antimicrobial Action of Compounds Containing 1-Oxo-2-methylene-, 1,2-Dioxo- and Derived Groupings" pages 345-351 * Pages 346, 347, compounds 3g, 5g, 6g, 3h, 5h, 6h and explanations; page 349, table 1 *		
X	--- CHEMICAL REVIEWS, vol. 61, 1961, American Chemical Society F.C.BROWN "4-Thiazolidinones" pages 463-521 * Page 480, table 5; page 509, right column, fourth passage; page 511, left column, lines 47-48 *	1-3, 23	
A	* Page 480, table 5 *	4, 10, 11, 14, 25	TECHNICAL FIELDS SEARCHED (Int. Cl.4)
D, A	--- JOURNAL OF THE ORGANIC CHEMISTRY OF THE USSR A TRANSLATION OF ZHURNAL ORGANICHESKOI KHIMII, vol. 14, no. 5, part 1, 1978 N.A.SHENBERG et al. "Investigation of the Reactivity and Tautomerism of Azolidines. XXIII. Alkylation of Salts of 5-Benzylidene-2-Thioxazolidin-4-ones" pages 1225-1228 * Page 1225, formulas Ia-If, IIIa-IIIIf; page 1226, compounds IIIa-IIIIf *	1, 2, 5, 16, 23, 25	
The present search report has been drawn up for all claims			
Place of search		Date of completion of the search	Examiner
VIENNA		15-11-1989	MAZZUCCO
CATEGORY OF CITED DOCUMENTS			
X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document		T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document	

THIS PAGE BLANK (USPTO)

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

THIS PAGE BLANK (02/10)

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☒ FADED TEXT OR DRAWING
- ☒ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☒ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.

THIS PAGE BLANK (USPTO)